

Occultation Newsletter

Volume III, Number 6

January, 1984

Occultation Newsletter is published by the International Occultation Timing Association. Editor and Composer: H. F. DaBoll; 6 N 106 White Oak Lane; St. Charles, IL 60174; U. S. A. Please send editorial matters, renewals, address changes, and reimbursement requests to the above, but for new memberships, new subscriptions, back issues, and any special requests, write to IOTA; P. O. Box 3392; Columbus, OH 43210-0392; U.S.A.

FROM THE PUBLISHER

For subscription purposes, this is the fourth and final issue of 1983.

O.N.'s price is \$1.40/issue, or \$5.50/year (4 issues) including first class surface mailing. Back issues through vol. 2, No. 13 still are priced at only \$1.00/issue; later issues @ \$1.40. Please see the masthead for the ordering address. Air mail shipment of O.N. back issues and subscriptions is 45¢/issue (\$1.80/year) extra, outside the U.S.A., Canada, and Mexico.

IOTA membership, subscription included, is \$11.00/year for residents of North America (including Mexico) and \$16.00/year for others, to cover costs of overseas air mail. European and U. K. observers should join IOTA/ES, sending DM 20.-- to Hans-J. Bode, Bartold-Knaust Str. 8, 3000 Hannover 91, German Federal Republic.

IOTA NEWS

David W. Dunham

The first IOTA annual meeting in Houston, TX, last November, was very successful, as reported by Charles Herold on p. 118. While in Texas, I also gave talks about last May's Pallas occultation at Rice University, Houston, and to a joint meeting of Ft. Worth and Dallas amateurs organized by George Ellis. George Ellis also attended the IOTA meeting, where he discussed possible cooperation of the Astronomical League with IOTA, especially for coordinating asteroid occultation observations. Don Archer already had sent me a listing of the addresses of all astronomical societies belonging to the Astronomical League, as well as of all (over 8000) individual A.L. members, conveniently arranged geographically by ZIP code. We noted how home computers might be used to disseminate last-minute predictions rapidly via mailgram-like services and computer bulletin boards.

Since IOTA must hold its annual meeting in Texas, Ellis suggested that we might hold the 1984 meeting in conjunction with the Texas Star Party in western Texas, the first weekend of June. This would result in a larger attendance and more publicity, as more amateur astronomers would be going there. However, observing the May 30th annular solar eclipse probably will make an early June meeting difficult, and it appears unlikely that three members of the IOTA board of directors, needed to conduct business,

would be able to attend. So the second annual meeting probably will be held late in the year again, in eastern Texas, and the 1985 meeting likely will be held with the Texas Star Party. However, we should not make too many plans until we hear from the Internal Revenue Service about our application for tax-exempt status. In late October, the I.R.S. sent us a letter asking for more information, and we have not yet received a reply to our response. If they deny tax-exempt status, IOTA probably will disincorporate.

I thank Rice University's Department of Space Physics and Astronomy for paying the cost of my round-trip plane ticket to Houston last November. The General Dynamics Recreation Association in Ft. Worth, Don Stockbauer, John Cotton, Paul Maley, and Walter Morgan provided lodging, meals, and other transportation.

The analysis of lunar occultation observations was set back severely last month when Tom Van Flandern left the U.S. Naval Observatory. Little progress with the analysis had been made during the last year as disputes with Captain Roberts and other administrators, about working hours and other matters, took more of Van Flandern's time. Although Captain Roberts prevailed, Van Flandern is undertaking legal actions to reinstate his position. In the meantime, the most sophisticated software for analysis of lunar occultation data sits idle in disk files at U.S. N.O. Since H.M. Nautical Almanac Office discontinued its lunar occultation work in 1981 (except for a low-level effort, which lasted another year, to complete some publications about its final analyses), the only workers I know who are paid to analyze lunar occultation data are now at the International Lunar Occultation Centre in Japan. Unfortunately, their manpower is limited; their work of preparing prediction information for publication, and of key-punching the observations and computing and distributing residuals, leaves little time for analysis. At U.S.N.O., Marie Lukac continues to work part time to produce the total occultation predictions, obtaining advice from Van Flandern and me. But the future of even this vital work is not certain. So far, Lukac and Alan Fiala have argued successfully for continuing the work, but there are some in the Nautical Almanac Office who would just as soon see it ended. But with Van Flandern not on the scene, occultations have lost a powerful spokesman. The finding that G is essentially constant will not help; see p. 118.

The ILOC finally distributed residuals for 1981 oc-

cultation timings to observers a few months ago. Some have been confused by the format of the distributed lists. I have begun an independent calculation of the residuals for a small sample of observations at U.S.N.O. and plan to publish the results in the next *O.N.*

On rare occasions, an SAO number less than 17 will appear in the USNO total occultation predictions. Since such SAO stars are north of declination +80°, they are not true SAO numbers. They are actually error codes for AGK3-catalog stars which are not in the SAO catalog.

In spite of late-night breakdowns of our van, both coming and going, we felt that the meeting of the American Astronomical Society's Division of Planetary Science in Ithaca, NY, last October, was very successful. R. Millis and E. Bowell described possible 1984 occultations by (1) Ceres and by Halley's Comet; see p. 129. Observations of occultations by (375) Ursula on 1982 November 15 and by (93) Minerva seven days later were described in *O.N.* 3 (3), 52. L. Wasserman reported in Ithaca that analyses of the observations at Lowell Observatory gave a diameter of 216 ± 4 km for Ursula and 173 ± 2 km for Minerva. The actual uncertainty is probably about ± 10 km due to lack of coverage near the limits of both occultations. Since no ellipticity was evident, only circular fits were made. S. J. Ostro reported that a double peak in a radar reflection with the Arecibo radio telescope for the small asteroid (2201) Oljato indicates that it might be a contact binary. Speckle observations of (532) Herculina and my first solution from analysis of last September's occultation by (51) Nemausa were reported at Ithaca and described in *Sky and Telescope* 67 (1), 60. I led a small meeting about publication of the results from the occultation of 1 Vulpeculae by (2) Pallas in May. We decided that the paper probably should be submitted to the *Astronomical Journal*. In addition to the occultation observations and analysis, we would list all miss observations to show the observational coverage in the vicinity of Pallas. We also would list everyone who travelled to observe the occultation, but were clouded out, to document the overall effort for this unique event. I hope to have a first draft of the paper ready in February. In the meantime, Paul Maley's account of the Texan effort for the event soon will be published in *Astronomy*, as well as translated versions in Italian, Latin American, and Swiss magazines. A first draft of the results from the occultation of 14 Piscium by (51) Nemausa should be ready in March or April.

Reports of lunar occultation timings made in the Soviet Union during 1980 and 1981 recently were received. Some grazing occultation observations were reported, including one of Aldebaran recorded at five stations on 1980 March 21 with the temperature -15°C! In a letter dated 1983 November 15, A. Osipov, Kiev, said that the Soviet report for 1982 was in press, and that a report for 1983 would follow in a few months. He reported that the occultation of 8.6-mag. SAO 104751 by (2) Pallas was observed from Engelhardt Observatory in Kazan on 1983 May 4, and that the occultation of 9.4-mag. SAO 41289 by (36) Atlante [Ed: misspelled Atalante in 1983 predictions] was seen in Khabarovsk last October 9th.

After the Soviet report for 1982 is received, and after the observations for 1982 are received on mag-

netic tape from ILOC, we will be able to make a final analysis of occultations timed during the total lunar eclipses which occurred that year. David Herald already has analyzed 105 timings during the July 6th eclipse and 141 timings during the December 30th eclipse made in Australia and New Zealand, and published the results in *Circular C83/9* of the Occultation Section of the Royal Astronomical Society of New Zealand (1983 October). In the same publication, Graham Blow reports that Peter Anderson timed a 6.7-second occultation of SAO 183401 by (10) Hygiea on October 8 at Bundaberg, Queensland. The disappearance was at 9h 47m 48s.3.

Hans-Joachim Bode recently sent me a report of the regional meeting held 1983 October 15 in Kiel, German Federal Republic. Asteroidal occultations were the main subject of discussion, and a cooperative network of observers of possible events in the G.F.R. and in Denmark was established. We hope that the network can be expanded to other European countries when the third European Symposium on Occultation Projects (ESOP III) will be held in June, probably on the 23rd, in either Czechoslovakia or Denmark. I will be attending an astrometry meeting near Munich earlier that week, so I plan to be at ESOP III.

The Astronomical League will be meeting in Milwaukee, WI, on 1984 August 1-4. I plan to be in the Los Angeles area at the time and will be unable to attend. Anyone who would be interested in speaking at a possible IOTA session or workshop during the meeting should contact Dan Koehler, W248 S7040, Sugar Maple Dr., Waukesha, WI 53186, phone 414,662-2987.

I apologize for my delays in preparing material for this issue; I know many of you are anxious to learn about this year's planetary and asteroidal occultations. Some information on early January events was distributed by Joseph Carroll in rough form along with the 1984 appulse predictions which he computes for all IOTA members. I again have not had time to publish organized accounts of grazes and asteroidal occultations and appulses, but plan to do so in the next issue, which will be targeted for late March.

THE TIMEKUBE/WEATHERADIO; WHETHER OR NOT

Don Stockbauer and Ken Thomson

[Ed: The following is an independent field evaluation, and solely represents the viewpoints of the authors.]

Many of us have waited eagerly for Radio Shack's newest Timecube/Weatheradio to appear in the stores, as a source of time signals. Stockbauer purchased one and was not pleased with the time signal reception at all! His old model Timecube almost always received WWV clearly on one or another of the three frequencies. Playing the two models simultaneously, he noticed that often the old one would receive a signal strongly while the new one would get only static on the same frequency. Sometimes the new one would receive a station other than WWV while the old unit picked up WWV quite well on that frequency. He asked another member of the Houston Astronomical Society, who is better versed in the mysteries of electronics, Thomson, to try to adjust the new unit for better reception. Thomson analyzed the whole design, and the results are quite discouraging!

He found that the new unit often receives an image frequency on top of WWV. For example, there often is a radioteletype signal transmitting at 15.910 MHz on the new unit; it is precisely at the predicted image frequency for a 455 KHz intermediate frequency (IF). It was found that both the old and new units have 455 KHz IF's by feeding a signal generator into the antenna. But the bad news is that while both receivers have two stages of IF amplification, the older Timecube has a stage of RF amplification, while the new one does not! As a result, we think the new Timecube will have much poorer RF selectivity, and nothing can be done about it, short of a redesign.

Stockbauer has kept his new toy to use as a 4th-level time backup on expeditions, and as an occasional weather source (the weather feature works fairly well).

We would appreciate feedback from other users of the new Timecube/Weatheradio, but would strongly advise having a backup for it, lest Murphy's law prevail. [Please route any replies via the editor, so that we may have the option of publishing them.]


GRAZING OCCULTATIONS

David W. Dunham

The grazing occultation predictions for the first half of 1984 either have been, or are being, sent to IOTA members. A correction that should be applied to the profiles for northern-limit waning-phase northern-latitude-libration grazes was described on p. 100 of the last issue. Contrary to what is stated there, no changes have been made to the ACLPPP for the 1984 predictions, so the described correction still should be applied when appropriate. It seems that many of the discrepancies noticed between the 78A and 80F prediction versions are due to the use of Perth 70 catalog data for the latter, even for bright northern-declination stars, which could not be observed well from Perth. The best solution to the current problems probably would be a new analysis of all the occultation data, using the XZ catalog as it stands for the southern stars, but removing the Perth 70 data from the northern stars. Unfortunately, Van Flandern's absence from USNO (see p. 113. guarantees that no such analysis will be done soon.

The southern-limit Cassini-region grazes also seem to be causing problems. Observations of a few grazes in these areas have shown large south shifts. Until better advice can be given from studying more observations, I recommend that observers apply a 0.5 southward correction to southern-limit ACLPPP profiles if more than half the profile points (including all within 1° of central graze) are 3's and 4's.

G. M. Appleby and L. V. Morrison recently published their "Analysis of lunar occultations - V. Grazing occultations 1964-1977" in *Mon. Not. R. Astr. Soc.* 205, p. 57. Observations of 800 grazes were studied to find corrections to Watts' charts, to the analytical lunar ephemeris



$j=2$, and to the zero point of right ascension in the FK4 catalog. Computer-drawn observation reduction profiles of 228 of the best-observed grazes were published on microfiche.

Robert Sandy, now in Independence, MO, has prepared some more reduction profiles for some recent grazes and some well-observed earlier events. These will be published as time and space permit in *O.N.*; some others will be distributed to expedition leaders.

Graham Blow photoelectrically recorded 3 disappearances and 2 reappearances during a near-graze of 8.5-mag. SAO 186844 at Mt. John Observatory in New Zealand last Sept. 15. Since the rotating buffer contained space for only 3.84 seconds of data, the star may have reappeared briefly again after the last-recorded disappearance. The predicted position angle was 16° from central graze.

Joan and I have rewritten and updated the papers describing the use of graze predictions, as part of a long-delayed larger general IOTA manual on occultations. Some other parts of the manual also have been prepared. These soon will be sent to Mark Allman for duplication and distribution to all new and current IOTA members. Allman also is updating some parts of *Observing Grazing Occultations VIII*, especially the useful section telling where to send requests and reports.


Again, time does not permit preparation of a list of observed grazes, but I am more determined than previously to include one in the next issue. After that, Don Stockbauer has agreed to collect reports of observations of grazes, make sure they are filled out properly, send blank forms when needed, and prepare the summaries for *O.N.* In the meantime, continue to send the reports to me, with, if possible, copies to ILOC.

LETTER

To the Editor:

Here are some notes relating to the article "Timing Occultations with a Digital Stop Watch," by Brian Loader (*O.N.* 3 (4), 85).

I have been using the described method for several years now, with only a slight difference; I have fixed the STOP key. In this way, I am sure of not losing the time base by accidentally stumbling or bumping, falling or grabbing the stop watch when I am late for a timing. In this way, I can use the time base for more than six months. I only un-fix the STOP key to reset the time base when it differs more than ten seconds from the correct time. For very



close events, I simply use a second stop watch (you can buy them for \$7 these days).


Some care must be taken in choosing the digital stop watch you will use. Some four years ago, I used the CASIO computer quartz CQ-1 to do my timings. When using the LAP key of that stop watch, you do influence the time base in some way. I found that when I used the LAP key some 100 times, the time base drifted about 0.2 second or more. Especially when you test the stop watch against the time signal to get an accurate difference, you could wind up with accuracy poorer than 0.1 second.

Henk Bulder
The Netherlands

LUNAR OCCULTATION OF ISEE-3

David W. Dunham

My orbital design work for the 3rd International Sun-Earth Explorer (ISEE-3) was mentioned briefly in *O.N.* 3 (3), 45. The complex orbit planned for the



spacecraft was shown on page 136 of last February's *Sky and Telescope*, and also in *Astronomy* 11 (9), 20 (1983 September). Co-workers at Computer Sciences Corp. and at Goddard Space Flight Center successfully executed the propulsive maneuvers needed to target the five lunar swingbys. The last swingby, on December 22nd, provided an unusual opportunity to combine my occultation work with the main part of my job. We planned to send ISEE-3 only 118 km above the mean lunar surface, much closer than the other swingbys, and the only one where the spacecraft would be occulted by the moon as seen from the earth. We predicted the time of occultation as seen from Goddard and Jet Propulsion Laboratory tracking stations in Alaska, Guam, and Australia. By comparing the observed with the predicted times, we hoped to estimate the actual height above the moon at closest approach.

I modified an existing program, which computes a topocentric ephemeris of a spacecraft, to calculate the angular distance to the center of the moon. When this distance was within 10" of the subtended lunar radius, I printed position angles and other quantities so that Watts' charts could be used to include limb corrections. The program was not completed and partially tested until Friday, Dec. 16. In the meantime, ISEE-3 was being tracked by radio at both Goddard and J.P.L. stations, and improved trajectories, determined from the tracking data, were given to us every two days. We needed the latest information in order to decide whether we needed to perform another propulsive maneuver on Dec. 20 in order to achieve the precise lunar swingby parameters needed to get close to Comet Giacobini-Zinner in 1985. In the meantime, an independent occultation prediction (without limb corrections, which changed the times by only 1 or 2 seconds) was being made with an older "coverage" program as a check, although it had not been designed for such a calculation. Comparison with the same trajectories was not made until Dec. 19, and they disagreed by 20 seconds! But computer hardware problems prevented making any program modifications needed to trace down the discrepancy; the machine still had not been fixed on Dec. 23. We looked for another independent Calculation. At J.P.L., the person who knew how to run their occultation program was on vacation. Marie Lukac, with some advice from Tom Van Flandern, finally overcame some computer problems at U.S.N.O. to compute a prediction only a few hours before the swingby. The U.S.N.O. times were closer to the "coverage" calculations, but disagreed with them by about ten seconds. However, the U.S.N.O. calculation neglected the spacecraft's motion, us-

ASTRONOMY DAY

MAY 5, 1984

For more information, contact:

Gary Tomlinson, Astronomy Day Coordinator
Astronomical League
Chaffee Planetarium
54 Jefferson S.E.
Grand Rapids, MI 49503
616-456-3985



The Astronomical Association of Northern California will be hosting Astronomy Day on 1984 April 7.

the spacecraft's motion, us-

ing fixed positions which I had supplied using the latest J.P.L. trajectory. The Goddard and J.P.L. trajectories obtained on Dec. 19 gave occultation times about seven seconds apart at a given station. At least, they indicated that the trajectory was accurate enough that the planned Dec. 20th correction maneuver could be cancelled.

As we waited for the swingby events in the control room at Goddard, we were confident that ISEE-3 would not crash into the moon, but wondered if it would survive a 29-minute passage through the moon's shadow. The spacecraft batteries had died a couple of years before, so sunlight was needed for the solar cells to provide all the spacecraft electrical power (in its originally designed "halo" orbit about the libration point 0.01 AU from the earth towards the sun, ISEE-3 was always in sunlight). Would hydrazine fuel which might still be in the lines freeze and expand, rupturing them and leaving ISEE-3 with no way to correct its orbit for the final targeting to the comet? Finally, over the intercom, we heard: "Alaska, LOS (loss of signal) at 17h 50m 46s G.M.T." I checked my sheet - only two seconds later than my predicted time using the Goddard trajectory, and one of those seconds would be explained by neglected light-time delay! LOS at Guam was 17h 52m 24s, four seconds late. The result for Ororral, Australia, was two seconds later than my time.

Emersion was at 18h 38m, but ISEE-3 was in eclipse then, and silent. Closest approach came six minutes later, over Mare Smythii. The eclipse ended at 18h 48m, after which we had to wait a few minutes for the spacecraft to power itself up. You could hear a pin drop as we waited and a command was sent to the spacecraft to report its condition. Then, "This is Guam - we have AOS (acquisition of signal)!" Robert Farquhar, who had conceived and led the effort to send ISEE-3 to Comet Giacobini-Zinner, blurted out, "Was there ever any doubt!" The readings from ISEE-3 showed that its temperature had dropped only 15° while in eclipse, and that all was well. A trajectory computed at J.P.L. using data both before and after the swingby showed that the closest approach was within 2 km of our target. The spacecraft, now leaving the earth-moon system, was rechristened the International Comet Explorer (ICE). We will be giving a series of papers about the trajectory design for this unique mission at the American Institute of Aeronautics and Astronautics' Astrodynamics Conference in Seattle, WA, this August. A 5-minute animation of ISEE-3's trajectory was videotaped at J.P.L.; I recently made a copy of it.

THE GRAVITATIONAL CONSTANT IS CONSTANT

David W. Dunham

In an article, "Experimental Test of the Variability of G Using Viking Lander Ranging Data," in *Physical Review Letters* 51 (18), 1983 October 31st issue, Hellings, R. W.; Adams, P. J.; Anderson, J. D.; Keese, M. S.; Lau, E. L.; and Standish, E. M., Jet Propulsion Laboratory; and Canuto, V. M. and Goldman, I., NASA Goddard Institute for Space Studies, NY, presented the result:

$$\dot{G}/G = (0.2 \pm 0.4) \times 10^{-11} \text{ per year.}$$

Using an interplanetary baseline for precise transponder range measurements gave these authors a big

advantage over previous efforts to measure G, as they state: "This (our) sensitivity represents more than an order of magnitude improvement over previous limits set with radar ranging data or with lunar orbit data. The quoted error is much larger than the formal standard deviation and represents uncertainties stemming from our lack of knowledge of the masses of the asteroids." In fact, the ranging measurements were so accurate that it was possible to determine the masses of the three largest asteroids, (1) Ceres, (2) Pallas, and (4) Vesta, as well as the mean densities of the 200 largest asteroids, broken down into C- and S-type groups. The result is consistent with cosmological theories, such as Einstein's, which predict no change in the gravitational constant, and rules out others, such as Dirac's large numbers hypothesis, which predict $\dot{G}/G =$ the inverse of the Hubble time of galactic expansion, or about 5×10^{-11} per year.

The "lunar orbit data" are primarily photoelectric timings of total occultations. Although a major breakthrough like the above is always welcome, it does remove a major rationale for lunar occultation timings. Of course, observations of total and grazing lunar occultations are still valuable for dynamical and astrometric studies (including the determination of proper motions around the sky, which in turn are needed for deriving the Oort parameters of Galactic rotation), resolution of close binary stars (many of which can not be detected by other techniques), photoelectric measurement of stellar angular diameters, and improvement of knowledge of the lunar profile (which is needed for analysis of solar eclipse observations used to determine small variations of the solar radius). With the virtual cessation of lunar laser ranging observations, occultation data again may become more important for determining lunar orbital parameters and Ephemeris Time.

MINUTES OF IOTA MEETING 1983 NOVEMBER 11

The first meeting of IOTA as a non-profit organization incorporated in the State of Texas was called to order on the 11th of November, 1983, at 7:15 P.M. It was a joint meeting with the Johnson Space Center Astronomical Society, held at the Lunar and Planetary Institute, on NASA Rd 1, Houston, Texas.

After some preliminary discussions of JSC Astronomical Society, Paul Maley introduced Dr. D. W. Dunham.

Dr. Dunham discussed an occultation of a star by the asteroid Minerva, in Europe and North America. He then presented some preliminary findings on the Pallas occultation. Of 130 sites in Texas, only 68 were able to escape the cloud cover and time the event. With all the chords presented, including some photoelectric observations, the profile of Pallas appears circular. Formal analysis still is to be completed. He also showed a video recording of a grazing occultation by the moon, that he and his wife made on 1983 October 27.

Paul Maley then talked about, and showed some 35 mm film slides of, the Indonesian eclipse expedition. He also showed some data that were collected by other IOTA members. He also discussed upcoming grazing occultations and expeditions planned by him and some other members through 1988.

Dr. Dunham explained some reasons for grazing occul-

tations:

1. for better astrometry (and refinements thereof);
2. for detection of double stars (only method suitable for some very close pairs);
3. for improved knowledge of the lunar position;
4. to find the north-south diameter of the moon more accurately; and
5. for the most accurate resolution of the lunar profile.

At 11:01 P.M. Dr. Dunham opened the IOTA portion of the meeting. There were 12 IOTA members present.

I. David Dunham made a motion to elect all candidates as presented on the ballot which was mailed to all members, as the officers of the organization. The motion was seconded by P. Maley. All members present voted in favor of the motion.

Charles Herold distributed the unopened envelopes with ballots from members to members present, who in turn, opened and counted the ballots. They then were collected by C. Herold, who totalled the groups of ballots. 132 ballots gave their voting rights, by proxy, to C. Herold. One ballot gave its voting rights to Mark Allman. C. Herold cast all votes for which he was proxy for the above motion. Mark Allman was not present.

The tally was 144 ayes (for), and 2 nays (against). The motion carried, as 59% of the members were in favor (30% is required in the bylaws).

II. A motion was presented to change membership expirations from a monthly to a quarterly basis. The motion was seconded by C. Herold.

There were 142 ayes (for), and 2 nays (against). The motion carried.

Dave Dunham mentioned that because of budget cuts at the U.S. Naval Observatory, it may become difficult to obtain grazing occultation predictions in the future (alternative means of obtaining predictions are being studied, in case such a catastrophe occurs).

George Ellis, of the Astronomical League (which has 8500 members), offered its services to IOTA, to help organize occultation observation events. He said that they have a communication network that is in operation all year long. This network could alert the most people who would be interested in becoming part of a grazing occultation expedition in the least amount of time.

The meeting was adjourned at 12:01 A.M., 1983 November 12.

Respectfully submitted

Charles H. Herold
Executive Secretary

MORE TELEPHONE TIME DEVELOPMENTS

David W. Dunham

A caution about a possible 0.25-second delay caused by the routing of many North American long-distance telephone calls through geosynchronous communications satellites was published on p. 106 of the last issue. Derald Nye informed me that WWV time signals are available within the U.S.A. on a toll-free num-

ber, 800,957-9999. Use of this number would have the same drawback as use of the other accurate telephone time numbers mentioned in the last issue. But the Naval Observatory has introduced a new number, 900,410-8463, which uses only land lines, not satellites. Hence, the time available via this number to all residents of the U.S.A. would be accurate to, at worst, a couple of hundredths of a second, less than visual timing uncertainties, and is consequently preferred over the other long-distance numbers for occultation work. The 900 numbers are not free, but cost 50 cents for each call, regardless of location within the U.S.A. U.S.N.O.'s earlier master clock time number, 653-1800, is still in operation, and can be used to obtain accurate time within the Washington, DC, free-calling area; presumably, the same is true for WWV's regular number in the Ft. Collins, CO, area. But others for whom these calls are long-distance should use the 900 number to avoid the possible satellite delay.

REVISED SOLAR RADIUS RESULTS

Marion Schmitz, Joan B. Dunham, and David W. Dunham

We recently discovered and corrected an error in our computer program which computes solar radius and lunar solar position corrections from reduced solar eclipse observations. Unfortunately, all of our previously published solar radius determinations must be revised, including those used in the article, "Solar Radius Change Between 1925 and 1979," published in *Nature* last August; see the end of IOTA News on p. 98 of the last issue of *O.N.* The new solar radius corrections for some eclipses are as follows:

Year	Correction
1715	+ 0.64 ± 0.2
1925	+ 0.08 ± 0.08
1979	- 0.11 ± 0.07
1980	- 0.03 ± 0.03

Hence, the change between 1925 and 1979 is - 0.19 or two times the standard error, not nearly as significant as the previous 4- σ result. The large decrease from 1715 to 1979 remains. Many other 19th- and 20th-century solar eclipse observations are in an advanced state of analysis.

PLANETARY OCCULTATIONS DURING 1984

David W. Dunham

Predictions of occultations of stars by major and minor planets during 1984 are given in two tables below. Reports of observations of these events should be sent to me at P. O. Box 7488, Silver Spring, Maryland 20907, U.S.A. (telephone 301, 585-0989), with (if possible; indicate on the form to whom copies are sent) a copy to Gordon Taylor, H. M. Nautical Almanac Office, Royal Greenwich Observatory, Herstmonceux Castle, Hailsham, Sussex BN27 1RP, England. Preferably, the report forms of the International Lunar Occultation Centre (ILOC), or the equivalent IOTA/ILOC graze report forms, should be used for reporting timed occultations or appulses (we want to know if you observed, and saw nothing, but do not need a report if weather prevented effective observation). The only difference from report-

(Text continues on page 121)

Table 1, Part A

U.T. UNIVERSAL P L A N E T	S	T	A	R	O C C U L T A T I O N	E1	M	0	0	N	Ephem.							
DATE	TIME	NAME	SAO No.	SP	Dec.	Am	Dur	df	P	Possible Area	SUN	E1	M	0	0	N	Up	Source
1983	Dec 30	20 ^h 29 ^m -33 ^m	Loreley	12.2 2.37	11.4	6 ^h 38 ^m 9	29°47'1.2	15°520	15	Mauritius, South Africa?n	173°140°	11-	none	Herget78				
1984	Jan 7	15 01-05 P/Halley	22.9 7.21	10.0 F8	6 24.9	10 18	12.9 3	16	42	Hawaii?s; Japan?s; southern USSR	180 119	17+	w 75°E	Yeomans7				
	Jan 7	22 23-38 Vesta	6.9 1.68	94222	9.1 K0	4 54.2	19 18	0.1 62	28	4 southern Africa, northeast Brazil	148 96	19+	w 5°W	APAEANAXX				
	Jan 12	7 42-45 P/Halley	22.9 7.19	12.3	6 21.9	10 23	10.6 3	16	104	northern Canada, Alaska	160 64	59+	w 90°W	Yeomans7				
	Jan 12	17 53-68 Bertha	13.0 2.99	12.0	2 07.4	20 20	1.3 34	57	22	Europe, northern Siberia?s	105 10	63+	all	EMP 1983				
	Jan 14	10 57-65 Pales	11.7 1.88	97945	9.3 K5	8 33.0	17 26	2.5 15	22	Tahiti, Samoa, Queensland	167 68	79+	w130°W	Herget77				
	Jan 16	8 06-18 Bertha	13.0 3.04	12.7	2 08.3	20 30	0.9 27	46	22	north Pacific, western Canada	102 48	93+	all	EMP 1983				
	Jan 20	7 00-09 Prokne	12.3 2.26	117122	9.3 K0	8 44.6	3 45	3.1 12	19	17 Brazil?s; GA-WA; Alaska?n	161 21	96-	all	EMP 1983				
	Jan 20	12 56-69 Loreley	12.4 2.43	78310	9.2 A2	6 20.4	29 01	3.3 18	24	15 Hawaii; Taiwan?n; Indochina	155 52	94-	e100°E	Herget78				
	Jan 20	14 55-69 Loreley	12.4 2.43	10.2 G0	6 20.3	29 01	2.4 18	24	15	HI?s; Japan?s; n. China, s.w.Asia	155 53	94-	e 65°E	Herget78				
	Jan 20	19 09-20 Ursula	12.4 2.45	10.2	8 38.3	31 35	2.3 14	20	17	Indonesia, Sri Lanka, cen. Africa	167 27	93-	all	EMP 1981				
	Jan 23	17 41-45 Hermione	12.8 2.84	10.5	2 34.5	12 46	2.4 15	25	21	northern Europe?s	98 153	66-	none	EMP 1983				
	Jan 24	16 01-13 Ursula	12.4 2.44	11.5	8 34.4	31 37	1.3 14	20	17	Hawaii, southern USSR, Mideast	168 83	56-	e120°E	EMP 1981				
	Jan 26	18 09-31 Daphne	12.6 2.46	112434A	8.8 A0	5 04.8	2 23	3.9 23	35	18 s.w. Australia, n. Africa, Spain	127 153	34-	e 95°E	Herget78				
	Jan 26	18 19-34 Daphne	12.6 2.46	112434B	11.8	5 04.8	2 23	1.3 23	35	18 Mauritius, e. cen. & n.w. Africa	127 153	34-	e 90°E	Herget78				
	Jan 27	2 01-06 P/Halley	22.9 7.19	14.2	6 13.1	10 40	8.7 3	17	104	Europe?s; Canada?s	145 144	30-	e 0°	Yeomans7				
	Jan 29	4 24-37 Hestia	13.0 2.09	118733	9.1 F5	11 11.3	3 42	3.9 17	36	23 s. Europe, U. K., n. Canada	141 98	13-	e 10°E	Herget77				
	Jan 29	4 56-60 Bertha	13.1 3.22	11.9	2 13.4	21 14	1.5 15	27	23	Alaska, Canada?s	90 132	13-	none	Herget78				
	Jan 29	22 20-26 Prokne	12.2 2.24	11.7	8 35.5	4 55	1.1 12	18	17	southern Africa?n; eastern Brazil	167 152	4-	none	EMP 1983				
	Jan 30	8 12-16 Bertha	13.2 3.26	10.9	2 14.6	21 23	2.4 14	25	23	e. Siberia, Alaska, w. Canada?s	88 107	3-	none	EMP 1983				
	Feb 3	21 27-40 Bamberg	10.9 1.93	11.6	9 14.2	20 09	0.4 17	18	11	w. Indonesia, India, n. Africa	176 160	3+	none	Herget77				
	Feb 6	20 37-54 Dembowska	10.7 2.21	10.8 K	11 16.1	15 21	0.7 13	27	22	n.w. Australia, northern Africa	151 154	20+	w 10°E	EMP 1975				
	Feb 10	2 03-06 Bertha	13.2 3.39	13.2	2 20.7	22 05	0.8 11	20	24	U. S. A.	80 13	49+	all	EMP 1983				
	Feb 13	16 06-08 Prokne	12.3 2.28	11.7	8 24.1	6 39	1.1 12	19	17	Japan?s; eastern Siberia	159 33	84+	all	EMP 1983				
	Feb 14	7 08-14 Bamberg	11.1 1.97	11.5	9 03.1	20 21	0.6 18	19	11	U. S. A.	168 26	89+	all	Herget77				
	Feb 15	0 41-58 Victoria	11.5 1.96	11.8	7 37.4	9 58	0.6 12	25	21	central Africa, southeast U.S.A.	148 15	94+	all	Herget77				
	Feb 17	12 28-29 Interamnia	12.3 3.65	186422	8.0 G5	18 08.1	-29 39	4.3 10	12	16 Mexico	56 117	99-	all	EMP 1977				
	Feb 17	16 35-48 Victoria	11.6 1.98	11.9	7 35.6	10 08	0.6 13	27	21	China, southwest U. S. S. R.	145 46	99-	all	Herget77				
	Feb 19	1 10-27 Metis	9.9 1.52	119464	9.3 F8	12 30.7	6 03	1.1 26	37	13 s.w.Asia,w.U.S.S.R.,s.Scandinavia	144 8	94-	all	Branham				
	Feb 19	16 55-57 Prokne	12.4 2.31	10.6	8 19.7	7 28	2.0 13	20	17	sothern Africa	153 64	89-	e 55°E	EMP 1983				
	Feb 20	16 34-37 Prokne	12.4 2.31	12.1	8 19.0	7 35	0.9 13	20	17	Mauritius, Kenya?	152 78	81-	e 70°E	EMP 1983				
	Feb 21	14 16-33 Victoria	11.7 2.01	11.9	7 33.2	10 21	0.6 15	31	22	New Zealand, Australia, India?n	140 102	72-	e120°E	Herget77				
	Feb 22	11 05-11 Adorea	12.1 1.81	11.0 K0	9 30.6	16 19	1.4 8	24	31	New Zealand,Australia (poor orbit)	167 87	64-	e160°E	EMP 1983				
	Feb 23	3 18-27 Vesta	7.8 2.17	11.4	4 52.6	21 33	0.0 54	28	6	northern South America	101 162	56-	e 60°W	APAEANAXX				
	Feb 23	4 15-18 Siegena	13.1 3.28	141366	9.0 G0	16 45.7	-3 26	4.2 9	17	23 Greenland; northwestern Europe?s	83 22	56-	all	EMP 1981				
	Feb 24	1 52-57 Vesta	7.8 2.18	11.9	4 53.2	21 36	0.0 52	27	6	central South America	100 174	46-	none	APAEANAXX				
	Feb 25	0 05-06 Euphrosyne	11.8 2.80	75140A	8.6 A0	2 03.4	24 52	3.2 9	11	15 southern South America?n	63 136	37-	none	EMP 1980				
	Feb 25	0 05-06 Euphrosyne	11.8 2.80	75140B	9.1	2 03.4	24 52	2.8 9	11	15 northern South America	63 136	37-	none	EMP 1980				
	Feb 25	16 25-39 Prokne	12.5 2.35	11.3	8 15.9	8 15.9	8 15.9	8 15.9	8 15.9	17 Australia, India, Mideast	147 142	30-	e125°E	EMP 1983				
	Feb 25	20 27-36 Metis	9.8 1.48	11.8	12 27.0	6 40	0.2 20	29	13	Japan, Siberia?s	151 88	29-	e120°E	Branham				
	Feb 26	11 18-27 Andromache	13.9 2.89	80634	7.7 G0	9 03.9	20 43	6.2 6	24	52 Samoa, southern Indonesia	156 146	24-	none	EMP 1983				
	Feb 28	15 18-24 Metis	9.7 1.47	11.8	12 25.1	6 56	0.1 19	27	13	Japan, China	154 122	9-	none	Branham				
	Mar 1	1 54-61 Prokne	12.5 2.39	11.3	8 13.6	8 49	1.5 15	24	18	northwestern Africa, Europe	142 156	3-	none	EMP 1983				
	Mar 2	18 42-45 Vesta	7.9 2.28	11.8	4 58.9	22 02	0.0 40	22	6	northern Europe, Siberia?s	94 94	0+	none	APAEANAXX				
	Mar 4	21 10-15 Germania	13.0 2.89	184336A	3.0 B2	16 18.1	-25 28	10.0 15	26	22 Tadzhik SSR; s.China, Philippines	97 119	4+	none	EMP 1981				
	Mar 4	21 08-13 Germania	13.0 2.89	184336B	5.2	16 18.1	-25 28	7.8 15	26	22 Kazak SSR, China, Taiwan?n	97 119	4+	none	EMP 1981				
	Mar 5	14 14-26 Elektra	12.8 2.99	12.2	13 54.1	11 49	1.1 16	24	18	New Zealand, Japan?s; e. Siberia	138 161	7+	none	EMP 1983				
	Mar 5	15 20-33 Bamberg	11.6 2.15	9.4	8 46.0	20 19	2.3 27	31	12	Manchuria, Mongolia, Kazakhstan	144 112	7+	w 75°E	Herget77				
	Mar 5	20 08-09 Euphrosyne	11.8 2.89	12.2	2 17.7	26 31	0.6 8	10	16	northwestern Africa	57 28	8+	w 5°E	Herget78				

Table 1, Part B

1984 UNIVERSAL P L A N E T	DATE	TIME	NAME	my	Δ , AU	S	T	A	R	Dec.	Δ m	O C C U L T A T I O N	P	Possible Area	E1	M	0	0	N	Up	Ephem.	Source		
				my		SAO No	my	Sp	R.A.	(1950)					SUN	E1	M	0	0	N	Up			
Mar 10	10 ^h 41 ^m 57 ^s		Thiabe	12.3	2.47		12.6		h ¹¹ 11 ^m 3	15° 41' 0.6	41° 55' 15	New Zealand?; Australia			132°	46°	47+			all		Herget78		
Mar 10	21 50-80	Victoria		12.0	2.20		12.1	7 27.6	11 21 0.7	40 86 24	South Africa?; Brazil?;s	121	32	52+ w 10°E	Herget77							Herget77		
Mar 13	5 53-66	Euterpe		9.8	1.21	118750		9.6	G5 11 13.0	8 03 0.8	12 23 15	Morocco, Labrador, northern Canada	173	52	76+ w 30°W	Herget77							Herget77	
Mar 14	3 48-57	P/Halley		23.0	7.52		13.8	5 55.2	12 00 9.2	9 48 109	Peru; Mexico?n	95	42	85+ all	Yeomans7								Yeomans7	
Mar 15	4 22-26	Metis		9.5	1.44		10.1	12 11.8	8 29 0.5	15 21 12	western Europe?;s	170	37	92+ w 10°E	Branham								Branham	
Mar 16	0 47-48	Junio		9.6	2.24		11.6	4 00.3	8 39 0.2	8 9 12	northeastern Canada	65	97	97+ all	APENAXX								APENAXX	
Mar 16	22 50-59	Prokne		12.8	2.55		12.3	8 08.6	10 42 1.0	22 36 19	Brazil?e	125	49	99+ all	EMP 1983								EMP 1983	
Mar 23	4 11-29	Kassandra		12.6	1.96	159989		8.5	A0 16 32.5	-16 10 4.1	21 44 22	Bermuda?; n.w. Africa, Italy	113	12	62- e 60°W	Herget78								Herget78
Mar 25	7 41-54	Saturn		1.0	9.08	158913		8.6	A5 14 53.9	-13 56 0.0	146m 46 0	Americas, Pacific	139	61	40- e125°W	NAO001								NAO001
Mar 26	11 18-19	Junio		9.7	2.35		11.0	4 21.0	10 13 0.3	75 8 13	western and central Australia	60	124	30- none	APENAXX									APENAXX
Mar 26	23 18-20	Hygiea		10.8	3.13	163443		9.3	G0 20 16.4	-19 44 1.8	15 11 10	southern India	64	6	25- all	Schmadel								Schmadel
Mar 27	9 21-23	Penelope		12.8	2.42	162170		8.1	B9 19 02.4	-16 57 4.7	4 13 38	northern Mexico, Cuba	81	27	22- e 90°W	EMP 1981								EMP 1981
Mar 31	19 18-21	Parthenope		10.2	1.63		11.5	G5 12 10.9	5 47 0.3	13 21 15	n.e. Siberia; Japan, China?;s	168	177	1- none	Herget77								Herget77	
Apr 2	17 41-57	Metis		9.7	1.48		12.3	11 54.2	9 53 0.1	17 24 13	southern Asia, eastern Europe	160	148	2+ none	Branham								Branham	
Apr 3	10 08-18	Bamberga		12.3	2.55		11.0	8 38.9	19 19 1.6	59 73 14	Australia	114	92	4+ w130°E	Herget77								Herget77	
Apr 9	23 06-10	Victoria		12.4	2.58		13.2	7 36.9	12 24 0.4	11 26 28	Newfoundland?; Scandinavia	94	14	59+ all	Herget77								Herget77	
Apr 10	4 51-55	Ursula		13.2	3.16		13.5	8 03.1	27 17 0.6	18 29 21	California, Mexico	97	6	61+ all	EMP 1981								EMP 1981	
Apr 10	6 49-61	Meliboea		12.9	2.36	138680		8.3	G5 12 13.3	-5 53 4.6	10 21 22	Argentina, Chile, Hawaii	165	62	62+ w 95°W	EMP 1982								EMP 1982
Apr 10	6 49-60	Meliboea		12.9	2.36	138680		10.3	12 13.3	-5 53 2.6	10 21 22	Patagonia; Hawaii?n	165	62	62+ w 95°W	EMP 1982								EMP 1982
Apr 13	1 23-28	Prokne		13.1	2.92		12.6	8 12.9	13 00 1.1	18 31 22	South America?n	100	44	89+ all	EMP 1983								EMP 1983	
Apr 14	5 00-02	Loreley		13.4	3.49		13.6	6 34.2	24 47 0.7	10 16 22	western North America?;s	74	84	96+ all	Herget78								Herget78	
Apr 19	1 01-03	Tamara		11.5	1.08	207516		8.8	16 14.2	-34 42 2.8	8 17 17	Seychelles, Madagascar; sAfrica?w	139	13	86- all	EMP 1983								EMP 1983
Apr 19	10 19-22	Prokne		13.2	3.01		9.7	F5 8 16.0	13 21 3.5	15 27 22	Japan; Alaska?n	94	133	83- e140°W	EMP 1983								EMP 1983	
Apr 21	23 25-26	Daphne		13.3	3.38		12.1	5 51.3	10 14 1.5	7 12 24	northeastern South America	57	152	60- none	Herget78								Herget78	
Apr 24	8 37-38	Vesta		8.3	2.92		11.0	6 05.7	24 15 0.1	17 10 8	Queensland	57	133	37- none	APENAXX								APENAXX	
Apr 24	21 37-38	Daphne		13.3	3.40		12.5	5 54.8	10 24 1.2	6 11 24	Brazil?;s	55	121	32- none	Herget78								Herget78	
Apr 27	14 56	Daphne		13.3	3.42		12.2	5 58.2	10 34 1.4	6 11 24	Kazakhstan	54	91	12- none	Herget78								Herget78	
Apr 30	21 22-23	Ursula		13.4	3.46		12.6	8 14.9	25 19 1.3	11 18 23	eastern Brazil	80	83	0- none	EMP 1981								EMP 1981	
May 2	1 27-40	Nemesis		12.0	2.13	139402		7.9	K0 13 30.4	-2 10 4.2	9 23 27	South Africa, Brazil, Peru	159	149	1+ none	Herget78							Herget78	
May 5	11 19-22	Siegna		12.3	2.39		12.0	16 59.0	4 48 0.9	16 25 17	New Zealand?n	142	143	19+ none	EMP 1981								EMP 1981	
May 8	2 29-31	Victoria		12.7	2.95		13.8	8 01.1	12 23 0.3	6 15 32	eastern U. S. A.	73	18	46+ all	Herget77								Herget77	
May 11	10 23-32	Athamantis		10.9	1.55	158162		8.5	K0 13 44.6	-16 50 2.5	11 23 19	New Zealand,Queensland?;NewGuinea	159	32	82+ all	EMP 1980							EMP 1980	
May 11	12 25-32	Tamara		11.0	0.96	226309		9.5	A0 15 48.7	-42 12 1.8	7 16 15	Indonesia	154	67	82+ all	EMP 1983							EMP 1983	
May 12	15 18-52	Mars		-1.9	0.54	159094		7.0	G5 15 09.6	-17 54 0.0	19m2 25 1	Hawaii, n. Australia, Asia	178	32	91+ w170°W	NAO001							NAO001	
May 12	21 01-02	Hektor		15.5	5.92		10.9	0 10.9	5 49 4.6	75 15 37	South China Sea; Australia?;s	47	162	93+ w105°E	EMP 1975								EMP 1975	
May 13	1 52-53	Loreley		13.6	3.86		13.1	7 05.2	23 17 1.0	7 12 25	southeastern North America?n	53	99	94+ all	Herget78								Herget78	
May 15	23 37-55	Tamara		11.0	0.95	226130		9.5	A0 15 41.0	-43 27 1.7	7 16 15	e. & s. Africa, Tierra de Fuego	155	23	99- all	EMP 1983								EMP 1983
May 16	9 43-52	Lutetia		10.8	1.47	158714		9.4	G5 14 37.3	-12 22 1.7	11 23 19	Guatemala; Mexico, Hawaii?n	165	30	98- all	EMP 1977								EMP 1977
May 16	18 26-28	Pallas		10.4	3.55		12.1	22 46.8	7 59 0.2	24 17 10	China, Japan	70	90	97- all	Sitarski								Sitarski	
May 18	4 46-68	Metis		10.7	1.91		11.0	11 38.8	8 52 0.6	28 45 16	s.w. U. S. A., Mexico, n. Chile	115	104	89- e113°W	Branham								Branham	
May 19	15 31-37	Athamantis		11.0	1.59	158099		9.2	F8 13 39.3	-15 40 2.0	13 28 20	Okinawa, n.e.China, cen. Siberia	150	85	78- s 40°N	EMP 1980							EMP 1980	
May 22	12 46-49	Fortuna		12.0	2.36		10.6	10 59.5	5 34 1.6	25 33 15	southeastern Siberia	103	164	52- none	EMP 1981								EMP 1981	
May 25	8 09-14	Fortuna		12.0	2.40		10.8	11 01.0	5 27 1.5	22 30 15	Fiji; Queensland, New Zealand?;s	101	161	26- none	EMP 1981								EMP 1981	
May 28	3 35-36	Thiabe		13.2	3.51		11.4	8 48.0	14 02 2.0	9 14 22	Mexico?n	64	94	7- none	Herget78								Herget78	
May 29	17 48-49	Diotima		13.1	3.44		14.0	9 08.8	26 55 0.4	7 13 24	Caucasus Mountains to N.W. India	64	75	1- none	EMP 1980								EMP 1980	
May 30	11 00-14	Klymene		13.6	2.64	186140		8.8	F8 17 59.3	-26 01 4.8	10 25 29	cen. Chile; N. Z., s. Australia?n	159	156	0- none	EMP 1983							EMP 1983	
May 31	10 04-05	Pallas		10.3	3.34		11.5	22 57.4	8 52 0.3	32 21 9	northern Chile, Bolivia	81	89	1+ none	Sitarski								Sitarski	
Jun 1	12 06-07	Io		13.5	3.39	117560		8.6	F8 9 14.5	7 27 4.9	6 14 33	southern Australia?n	68	47	4+ none	Herget78							Herget78	
Jun 2	0 18-19	Aglaja		12.6	2.35	146574		9.1	K0 23 11.1	-9 04 3.6	8 15 22	South Africa?n	86	115	6+ none	Herget78							Herget78	
Jun 2	12 21-30	Tamara		11.1	0.96	225520		9.8	15 08.3	-46 55 1.6	10 20 15	Indonesia, Singapore	149	127	9+ w120°E	EMP 1983							EMP 1983	
Jun 2	17 36-49	Marianna		13.1	2.29	227909		8.5	F8 17 20.5	-44 05 4.6	11 23 24	Australia, southern Africa	157	148	11+ w 35°E	Herget78							Herget78	

are presented in the same format as those for last year's events. Due to the large number of events, the tables are given on alternating pages, so that all the data for a given event (and for others which occur either just before and/or just after it) are available on facing pages. Explanations of the data

given in the tables, and of the finder charts, and regional and world maps appearing in *O.N.*, as well as information about local circumstances (appulse predictions) sent to IOTA members were given in the article about 1983 events in *O.N.* 3 (1), 9. Joseph E. Carroll, 4261 Queen's Way, Minnetonka, MN 55345, (Text continues on page 125)

Table 2, Part B

1984 DATE	M I N O R No.	Name	P L A N E T km-diam.-"	R S O I Type	MOTION °/Day	S T A R SAO No	A R DM No.	D	S T E L L A R m'	m	Time	df	S	A G K 3 No	Shift	A P P A R E N T R.A.	Dec.	
Mar 10	88	Thisbe	232 0.13	1465 C	0.076 285°								C			8 ^h 13.2 ^m	15°35'	
Mar 10	12	Victoria	135 0.08	575 S	0.051 344								C			7 29.5	11 17	
Mar 13	27	Euterpe	116 0.13	355 S	0.253 293	118750	+08°2478	0.15	136	15	0.6	AX	N 8°1465	0.14	0.9	11 14.8	7 52	
Mar 14		P/Halley	100 0.02	988	0.050 307								C			5 57.1	12 00	
Mar 15	9	Metis	168 0.16	679 S	0.254 292								C			12 13.6	8 17	
Mar 16	3	Junio	267 0.16	1144 S	0.507 72								C			4 02.2	8 45	
Mar 16	194	Prokne	195 0.11	1134 C	0.113 343								C			8 10.5	10 36	
Mar 23	114	Kassandra	131 0.09	488 C	0.105 72	159989	-15 4346	0.07	106	17	0.4	PX		-0.00	0.9	16 34.4	-16 14	
Mar 25		Saturn	115644 8.78		0.047 290	158913	-13 4022	0.08	538	41	0.9	X				14 55.8	-14 05	
Mar 26	3	Junio	267 0.16	1153 S	0.516 74								C			4 22.9	10 18	
Mar 26	10	Hygiea	443 0.20	3414 C	0.322 75	163443	-20 5887	0.09	198	6	0.6	X				20 18.4	-19 38	
Mar 27	201	Penelope	92 0.05	281 M	0.329 80	162170	-17 5459	0.08	135	6	0.4	X				19 04.3	-16 54	
Mar 31	11	Parthenope	155 0.13	649 S	0.240 295		+ 6 2571	0.06	76	6	0.3	A	N 5 1701			12 12.6	5 36	
Apr 2	9	Metis	168 0.16	686 S	0.220 283								C			11 56.0	9 41	
Apr 3	324	Bamberga	256 0.14	1629 C	0.056 152								C			8 40.9	19 11	
Apr 9	12	Victoria	135 0.07	572 S	0.156 83								C			7 38.7	12 19	
Apr 10	375	Ursula	214 0.09	1385 C	0.128 136								C			8 05.2	27 11	
Apr 10	137	Meliboea	153 0.09	814 C	0.206 309	138680	-05 3463 A	0.28	481	33	1.6	X				12 15.1	-6 04	
Apr 10	137	Meliboea	153 0.09	814 C	0.206 309	138680	-05 3463 B						X			12 15.1	-6 04	
Apr 13	194	Prokne	195 0.09	1136 C	0.122 60								C			8 14.8	12 53	
Apr 14	165	Loreley	228 0.09	1486 C	0.217 103								C			6 36.2	24 46	
Apr 19	326	Tamara	90 0.12	215 C	0.365 198	207516	-3410911 B						G			16 16.4	-34 47	
Apr 19	194	Prokne	195 0.09	1136 C	0.141 69		+13 1887	0.08	170	13	0.5	A	N13 828			8 17.9	13 15	
Apr 21	41	Daphne	204 0.08	1109 C	0.303 78								C			5 53.1	10 14	
Apr 24	4	Vesta	555 0.26	4248 U	0.365 87								C			6 07.7	24 15	
Apr 24	41	Daphne	204 0.08	1106 C	0.309 79								C			5 56.7	10 25	
Apr 27	41	Daphne	204 0.08	1104 C	0.315 79								C			6 00.1	10 34	
Apr 30	375	Ursula	214 0.09	1386 C	0.191 121								C			8 16.9	25 12	
May 2	128	Nemesis	116 0.08	498 CEU	0.194 282	139402	-01 2834 A	0.55	849	68	3.0	AG	S 2	809	0.66	1.4	13 32.1	-2 21
May 5	386	Siegna	203 0.12	1207 C	0.172 310								C			17 00.7	4 45	
May 8	12	Victoria	135 0.06	569 S	0.258 94								C			8 03.0	12 17	
May 11	230	Athamantis	116 0.10	406 S	0.226 310	158162	-16 3744	0.41	461	43	1.9	X				13 46.4	-17 00	
May 11	326	Tamara	90 0.13	212 C	0.426 226	226309	-41 7369	0.04	27	2	0.1	S				15 51.1	-42 18	
May 12		Mars	6782 17.46	*****	0.364 278	159094	-17 4273	0.51	199	34	1.4	YG		1.05	-0.5	15 11.6	-18 02	
May 12	624	Hektor	234 0.05	2437 D	0.184 57								C			0 12.6	6 00	
May 13	165	Loreley	228 0.08	1485 C	0.279 102								C			7 07.2	23 13	
May 15	326	Tamara	90 0.13	212 C	0.421 231	226130	-43 7243	0.04	27	2	0.1	S				15 43.3	-43 34	
May 16	21	Lutetia	114 0.11	386 M	0.235 282	158714	-11 3783	0.17	183	18	0.8	X				14 39.1	-12 31	
May 16	2	Pallas	538 0.21	5374 U	0.209 71								C			22 48.6	8 10	
May 18	9	Metis	168 0.12	702 S	0.104 145								C			11 40.6	8 41	
May 19	230	Athamantis	116 0.10	405 S	0.187 316	158099	-15 3723	0.11	127	14	0.5	X				13 41.2	-15 51	
May 22	19	Fortuna	226 0.13	1211 C	0.129 108								C			11 01.3	5 23	
May 25	19	Fortuna	226 0.13	1212 C	0.143 109								C			11 02.8	5 16	
May 28	88	Thisbe	232 0.09	1456 C	0.255 104								C			8 49.9	13 54	
May 29	423	Diotima	209 0.08	1219 C	0.277 114								C			9 10.8	26 47	
May 30	104	Klymene	134 0.07	721 C	0.161 266	186140	-2612698	0.13	246	19	0.8	X				18 01.4	-26 01	
May 31	2	Pallas	538 0.22	5351 U	0.166 72								C			22 59.1	9 03	
Jun 1	85	Io	149 0.06	739 C	0.249 95	117560	+07 2102	0.15	357	14	1.0	AS	N 7 1315	-1.01	-1.3	9 16.3	7 19	
Jun 2	47	Aglaia	156 0.09	627 C	0.288 67	146574	-09 6147	0.31	538	26	1.8	X				23 12.9	-8 53	
Jun 2	326	Tamara	90 0.13	210 C	0.326 247	225520	-46 7250						S			15 10.7	-47 03	
Jun 2	602	Marianna	139 0.08	685 C	0.189 270	227909	-4411668	0.18	295	23	1.0	YG			-1.77	9.2	17 23.0	-44 07

Table 1, Part C

1984 DATE	UNIVERSAL TIME	P L NAME	A NAME	N E T my	S AU Δ	T SAO No	S No	A R Sp	R A. (1950) Dec.	O C C U L T A T I O N Dur df P Possible Area	E1 SUN	M El	O %Sn1	O Up	N Up	Ephem. Source
Jun 2	19 22 ^m	24 ^m Hektor	15.4	5.65	10.7	65	10.7	65	0 ^h 22 ^m 6	7° 49' 4.7	9° 18' 35	Australia	64° 102°	11+	none	EMP 1975
Jun 4	0 56	Panopaea	10.9	1.28	211413	10.5	A0	19	33.7	-36 04	1.0	24	35	12	none	Hergel77
Jun 4	5 48-57	Tamara	11.1	0.96	225478	9.3	A0	15	05.3	-47 08	2.0	10	21	15	none	EMP 1983
Jun 4	18 29-30	Eugenia	12.9	3.32		11.2		1	17.2	4 37	1.9	7	11	19	none	Hergel77
Jun 8	18 23-24	Egeria	11.8	2.70	80995	9.2	G0	9	41.8	27 31	2.7	7	10	16	none	Hergel77
Jun 14	16 31-51	Dido	12.8	2.12	189495	9.1		20	34.4	-28 27	3.7	26	54	22	all	EMP 1983
Jun 20	16 58-60	Hektor	15.3	5.39		12.8		0	30.2	9 21	2.6	12	24	33	all	EMP 1975
Jun 23	20 14-15	Iris	10.0	2.26		12.9		2	04.3	17 18	0.1	6	8	15	all	Branham
Jun 23	22 30-31	Thise	13.2	3.80		12.2		9	17.8	11 59	1.4	7	11	24	none	Hergel78
Jun 24	4 25-32	Adeona	12.6	2.00	159236	9.0	G5	15	21.5	-14 31	3.6	19	38	21	none	ITA 1977
Jun 24	13 08-12	Fortuna	12.4	2.83	118864	6.4	K0	11	24.2	3 17	6.0	10	15	18	none	EMP 1981
Jun 27	13 21	Iris	9.9	2.22		13.3		2	12.4	18 01	0.0	6	8	15	none	Branham
Jun 29	1 13-15	Hektor	15.3	5.26		11.3		0	32.8	9 59	4.0	15	29	33	none	EMP 1975
Jun 30	6 14-23	Dido	12.6	2.02	189347	8.9	K0	20	26.7	-29 12	3.7	15	30	21	none	EMP 1983
Jul 6	5 54	Ceres	9.2	3.21		12.5		2	43.5	7 19	0.0	35	12	5	none	APAXAX
Jul 6	9 18-19	Hebe	10.1	2.45		11.2		3	43.7	8 13	0.3	5	8	19	none	Branham
Jul 7	18 38-44	Elektra	13.4	3.71		12.0		13	13.3	14 52	1.7	15	24	23	all	EMP 1983
Jul 7	18 42	Hebe	10.1	2.44	111439	8.1	G5	3	46.8	8 19	2.1	5	8	19	none	Branham
Jul 13	19 41-45	Cybele	12.8	3.26		12.7		1	14.9	6 06	0.8	22	25	15	all	Hergel78
Jul 14	1 01-04	Juewa	12.0	1.93	209985	8.3	G5	18	15.9	-39 42	3.7	15	25	17	all	Hergel78
Jul 15	9 20	Iris	9.8	2.05		11.2		2	51.9	21 09	0.3	7	8	13	all	Branham
Jul 16	0 34-53	Siegna	12.4	2.48		11.4		K0	16	13.6	5	19	14	23	all	EMP 1981
Jul 16	21 02-17	Isolda	12.7	2.33	164173	9.1	F5	21	06.3	-12 31	3.7	14	26	20	all	Hergel81
Jul 18	7 37-57	Lydia	11.3	1.58	211065	9.2	K0	19	10.5	-30 17	2.2	10	25	22	all	Hergel74
Jul 22	6 48-56	Neptune	7.9	29.38	186001	8.7	K5	17	54.7	-22 14	0.4	42 ^m	54	1	none	JPL DE96
Jul 28	10 45	Aurora	12.3	2.26	211730	9.6		19	57.7	-31 19	2.7	15	23	17	none	EMP 1981
Jul 30	10 13	Winchester	11.8	2.22		11.4		4	23.9	5 16	1.0	6	8	15	none	EMP 1982
Aug 2	6 32-33	Iris	9.7	1.88		10.0		3	31.6	23 40	0.6	8	9	12	none	Branham
Aug 2	16 46-55	Pallas	9.3	2.43		11.7		23	07.1	7 54	0.1	49	28	7	none	Sitarski
Aug 4	10 06-09	Tamara	12.2	1.35	225490	9.4	G5	15	05.6	-48 28	2.9	8	21	22	all	EMP 1983
Aug 5	23 52-53	Iris	9.6	1.84		11.4		3	39.8	24 07	0.2	8	9	12	none	Branham
Aug 7	9 24-26	Iris	9.6	1.83		11.7		3	42.8	24 16	0.2	8	9	12	none	Branham
Aug 8	17 43-59	Sylvia	11.7	2.29	211985	10.0	G2	20	15.3	-32 15	1.9	24	26	12	all	Hergel78
Aug 17	3 48-52	Ceres	8.8	2.64		11.8		3	25.1	9 36	0.1	70	22	4	all	APAXAX
Aug 19	13 40	Winchester	11.6	2.05		11.5		5	05.9	5 23	0.8	7	9	14	all	EMP 1982
Aug 19	20 49-50	Hebe	9.9	2.13		10.1		5	16.0	9 12	0.7	6	10	17	all	Branham
Aug 19	21 16-20	Pallas	9.0	2.28		12.0		22	57.7	5 27	0.1	33	18	6	all	Sitarski
Aug 20	9 39-40	P/Halley	22.5	7.07		14.0		6	39.6	13 41	8.5	6	31	102	all	Yeomans7
Aug 21	22 36-37	Winchester	11.6	2.03		11.9		5	10.6	5 21	0.6	7	9	14	all	EMP 1982
Aug 23	5 06-08	Ceres	8.7	2.56		11.7		3	29.3	9 44	0.1	83	26	4	all	APAXAX
Aug 25	9 00-19	Klymene	14.1	3.10	185260	9.2	K0	17	16.0	-25 40	4.9	27	69	34	none	EMP 1983
Aug 28	5 48-78	Eugenia	12.2	2.30		12.7		2	28.5	7 45	0.5	62	74	13	none	Hergel77
Aug 28	17 08-09	Winchester	11.6	1.97		12.6		5	23.8	5 11	0.4	7	10	14	none	EMP 1982
Aug 30	15 57	Winchester	11.6	1.95		11.7		5	27.5	5 08	0.7	8	10	14	none	Hergel78
Sep 1	0 09-18	Doris	12.0	2.42		12.2		2	45.0	12 49	0.6	41	64	18	none	Hergel77
Sep 1	1 19-32	Hektor	14.8	4.41		13.6		0	27.5	12 38	1.5	17	30	27	none	EMP 1975
Sep 2	10 12-13	Winchester	11.5	1.93		10.3	A0	5	32.6	5 02	1.5	8	10	13	none	EMP 1982
Sep 4	0 45-90	Dido	13.0	2.24	188498	9.3	K	19	41.1	-29 16	3.7	33	71	24	none	EMP 1983
Sep 5	21 09-12	P/Halley	22.4	6.70		10.2		6	44.0	13 23	12.2	9	47	97	all	Yeomans7
Sep 14	4 10-12	Hebe	9.8	1.91		10.7		5	59.7	8 03	0.4	8	12	15	all	Branham
Sep 14	21 22-24	Hebe	9.8	1.91		12.0		6	00.8	8 01	0.1	8	12	15	all	Branham

computes the appulse predictions. Special requests for the appulse predictions should be sent to his address, which was given incorrectly in the O.N. article about 1983 events, and also in the article about 1984 planetary occultations in this month's issue of *Sky and Telescope*. Specific information

about some of the events is given at the end of this article.

Sources of Planetary Occultations. The total of 230 events given in the list is the largest number ever published for a given year (actually, one event not
(Text continues on page 127)

Table 2, Part C

1984 DATE	M I N O R No. Name	P L A N E T km-diam.-"	R S O I Type	M O T I O N °/Day	S T A R SAO No	T A R DM No.	A R D	S T E L L A R D m"	D I A M E T E R m	T I M E Time	d f	S	C O M P A R I S O N AGK3 No	D A T A Shift Time	A P P A R E N T R.A.	D E C.
Jun 2	624 Hektor	234 0.06	2436 D	0.152 53°	191 211413	+ 7°	46	0.09	381	15	0.8	A	N 7°	45	h ²⁴ m ⁴	8°00'
Jun 4	70 Panopaea	153 0.16	529 C	0.166 191	211413	-3613626		0.02	23	4	0.1	S			19 36.0	-35 59
Jun 4	326 Tamara	90 0.13	210 C	0.311 248	225478	-46 7222		0.04	30	3	0.2	S			15 07.7	-47 16
Jun 4	45 Eugenia	250 0.10	1452 U	0.333 73											1 18.9	4 48
Jun 8	13 Egeria	245 0.13	1180 C	0.413 121	80995	+27 1804		0.12	238	7	0.7	AS	n27 1026	-1°16	9 43.8	27 22
Jun 14	209 Dido	137 0.09	611 C	0.082 237	189495	-2816800		0.08	237			S			20 36.5	-28 20
Jun 20	624 Hektor	234 0.06	2436 D	0.118 47								C			0 32.0	9 32
Jun 23	7 Iris	222 0.14	813 S	0.559 70								C			2 06.1	17 28
Jun 23	88 Thisbe	232 0.08	1449 C	0.304 107								C			9 19.6	11 51
Jun 24	145 Adeona	137 0.09	589 C	0.122 245	159236	-14 4194		0.20	296	40	1.1	X			15 23.4	-14 38
Jun 24	19 Fortuna	226 0.11	1222 C	0.257 111	118864	+03 2502	A	0.65	1331	61	4.0	ZA	N 3 1496	0.28	0.2 11 25.9	3 06
Jun 27	7 Iris	222 0.14	811 S	0.557 70								C			2 14.3	18 11
Jun 29	624 Hektor	234 0.06	2436 D	0.100 42								C			0 34.6	10 10
Jun 30	209 Dido	137 0.09	613 C	0.154 252	189347	-2917110		0.25	367	39	1.3	S			20 28.8	-29 05
Jul 6	1 Ceres	1025 0.44	12255 C	0.305 75								C			2 45.3	7 28
Jul 6	6 Hebe	186 0.10	637 S	0.549 83								C			3 45.6	8 20
Jul 7	130 Elektra	235 0.09	1754 C	0.143 138								C			13 15.0	14 41
Jul 7	6 Hebe	186 0.11	637 S	0.548 83	111439	+08 573		0.30	538	13	1.7	AS	N 8 398	-0.95	0.2 3 48.6	8 25
Jul 13	65 Cybele	311 0.13	2428 C	0.142 74								C			1 16.7	6 16
Jul 14	139 Juewa	165 0.12	788 C	0.188 283	209985	-3912452		0.15	211	19	0.8	PS		-1.26	-0.1 18 18.3	-39 41
Jul 15	7 Iris	222 0.15	801 S	0.543 73								C			2 53.9	21 17
Jul 16	386 Siegena	203 0.11	1174 C	0.115 211		+ 5 3174		0.09	156	18	0.5	A	N 5 2112		16 15.3	5 14
Jul 16	211 Isolda	168 0.10	925 C	0.166 260	164173	-12 5915		0.10	173	15	0.6	X			21 08.2	-12 23
Jul 18	110 Lydia	102 0.09	342 U	0.209 259	211065	-3016780		0.24	273	27	1.1	S			19 12.7	-30 14
Jul 22	Neptune	50184 2.36	*****	0.022 269	186001	-2212373		0.59	12547	633	11.7	HX		0.35	12.5 17 56.8	-22 14
Jul 28	94 Aurora	191 0.12	1106 C	0.189 266	211730	-3117238						S			19 59.9	-31 14
Jul 30	747 Winchester	208 0.13	764 C	0.536 87								C			4 25.7	5 21
Aug 2	7 Iris	222 0.16	793 S	0.519 76								C			3 33.6	23 47
Aug 2	2 Pallas	538 0.31	5229 U	0.150 222								C			23 08.9	8 05
Aug 4	326 Tamara	90 0.09	208 C	0.267 92	225490	-48 7147		0.07	64	6	0.3	S			15 08.0	-48 36
Aug 5	7 Iris	222 0.17	791 S	0.512 77								C			3 41.8	24 13
Aug 7	7 Iris	222 0.17	791 S	0.510 77								C			3 44.9	24 23
Aug 8	87 Sylvia	275 0.17	1913 P	0.163 254	211985	-3215863		0.09	150	13	0.5	S			20 17.5	-32 08
Aug 17	1 Ceres	1025 0.53	12140 C	0.184 81								C			3 27.0	9 43
Aug 19	747 Winchester	208 0.14	762 C	0.499 92								C			5 07.7	5 26
Aug 19	6 Hebe	186 0.12	650 S	0.467 93								C			5 17.9	9 14
Aug 19	2 Pallas	538 0.33	5190 U	0.239 224								C			22 59.5	5 38
Aug 20	P/Halley	100 0.02	829	0.078 101								C			6 41.5	13 39
Aug 21	747 Winchester	208 0.14	762 C	0.493 92								C			5 12.5	5 24
Aug 23	1 Ceres	1025 0.55	12122 C	0.159 83								C			3 31.2	9 51
Aug 25	104 Klymene	134 0.06	710 C	0.053 87	185260	-2512079		0.30	683	136	2.0	X			17 18.1	-25 42
Aug 28	45 Eugenia	250 0.15	1478 U	0.058 134								C			2 30.4	7 54
Aug 28	747 Winchester	208 0.15	762 C	0.475 94								C			5 25.6	5 13
Aug 30	747 Winchester	208 0.15	762 C	0.470 94								C			5 29.3	5 10
Sep 1	48 Doris	200 0.11	1092 C	0.066 110								C			2 46.9	12 58
Sep 1	624 Hektor	234 0.07	2434 D	0.105 271								C			0 29.3	12 50
Sep 2	747 Winchester	208 0.15	763 C	0.461 95		+ 4 978		0.03	37	1	0.1	A	N 5 640		5 34.5	5 03
Sep 4	209 Dido	137 0.08	619 C	0.062 325	188498	-2916476		0.29	469	112	1.6	S			19 43.3	-29 11
Sep 5	P/Halley	100 0.02	811	0.054 111								C			6 45.9	13 21
Sep 14	6 Hebe	186 0.13	662 S	0.386 100								C			6 01.6	8 04
Sep 14	6 Hebe	186 0.13	662 S	0.383 100								C			6 02.7	8 01

Table 1, Part D

1984 UNIVERSAL P L A N E T										O C C U L T A T I O N										Ephem.		
DATE	TIME	NAME	my	Δ _{AU}	S	T	A	R	Dec.	Δm	Dur	df	P	Possible Area	SUN	EI	M	O	0	N	Source	
					Sp	R.A.(1950)	h ^m	°01'	8°01'	0.2	8 ^s 12	15	northwest Siberia	82°	49 ^c	82-	all	Branham				
Sep 14	21	55 ^m -56 ^m Hebe	11.7	9.8 1.91	11.7	6 ^h 00 ^m 9	6 01.7	7 58 0.3	8 12 15	southwest Canada	82	44	78-	all	Branham							
Sep 15	11	01-03 Hebe	10.9	9.8 1.90	10.9	6 02.2	7 57 0.1	9 12 15	China, Japan	82	40	75-	all	Branham								
Sep 15	18	21-22 Hebe	12.3	9.8 1.90	12.3	6 02.2	7 57 0.1	9 12 15	China, Japan	82	40	75-	all	Branham								
Sep 16	2	18-26 Aglaja	8.9	11.3 1.54	146599	8.9	K0 23 13.4	-7 10 2.5	16 25 14	Iceland, southern Canada	173	70	72-	e 90°W	Hergert78							
Sep 17	14	26-29 Iris	12.2	9.1 1.43	12.2	5 05.2	26 53 0.1	14 14 9	eastern Siberia	97	4	58-	all	Branham								
Sep 17	18	53-85 Flora	10.2	8.4 0.96	10.2	60	1 51.1	-0 47 0.2	43 52 9	e. China, s.e. Asia	147	53	56-	e 70°E	Branham							
Sep 21	2	54-57 Winchester	11.4	1.77	13.0	6 04.6	4 06 0.2	10 12 12	central Africa	87	39	22-	e 10°W	EMP 1982								
Sep 23	8	52-56 Winchester	11.3	1.75	11.0	6 08.1	3 58 0.9	10 13 12	Central America, n. South America	88	66	5-	e 70°W	EMP 1982								
Sep 23	17	30-32 Winchester	11.3	1.75	11.9	6 08.6	3 56 0.5	10 13 12	eastern China, Japan	88	71	3-	none	EMP 1982								
Sep 23	19	40-50 Faina	12.4	1.72	11.1	5 02.4	12 49 1.6	11 26 22	Sri Lanka, southeast Asia, Taiwan	104	87	3-	e130°E	EMP 1982								
Sep 25	6	38-42 Iris	9.4	9.0 1.36	77119	9.4	A0 5 17.7	26 58 0.6	16 16 9	central South America	101	103	0+	none	Branham							
Oct 2	1	02-24 Chicago	13.2	3.01	164723	9.0	K2 21 51.1	-15 40 4.2	34 58 22	nw Africa, Canarlys., n.S.America	136	44	52+	w 25°W	EMP 1983							
Oct 6	3	04-15 Corduba	13.1	1.78	145486	8.6	K0 21 31.0	-6 36 4.6	17 42 24	western NorthAmerica (poor ephem)	130	15	87+	all	EMP 1982							
Oct 6	8	25-27 Hebe	11.0	9.6 1.72	11.0	6 29.1	6 23 0.3	13 18 13	Canada	95	121	88+	w105°W	Branham								
Oct 15	13-14	Penelope	12.0	1.83	162800	9.2	K2 19 33.0	-19 30 2.9	6 18 29	West and South Australia	99	45	90+	all	EMP 1981							
Oct 9	0	04-06 Iris	12.1	8.7 1.24	12.1	5 36.0	26 50 0.0	23 23 8	South Africa?n	111	80	99+	all	Branham								
Oct 9	3	45-58 Iris	11.0	8.7 1.24	11.0	5 36.2	26 50 0.1	24 23 8	Amazon, northwest Africa	111	78	99+	all	Branham								
Oct 9	15	16-19 Hebe	9.7	9.5 1.69	9.7	6 32.5	6 07 0.7	14 19 13	cen. & e.Australia, n. New Zealand	97	85	100+	all	Branham								
Oct 9	23	50-69 Doris	11.1	11.3 2.04	11.1	2 38.3	10 32 0.9	20 28 15	Virginia, Bermuda, central Africa	155	24	100-	all	Hergert77								
Oct 12	11	20-30 Hispania	11.3	1.66	74936	8.7	G5 1 44.9	28 14 2.7	15 22 14	Hawaii?n; eastern Indonesia	158	21	94-	e120°E	Hergert							
Oct 13	22	59-62 Winchester	11.1	1.57	10.9	6 34.8	2 36 0.9	16 19 11	northern Europe?n; n.w. Siberia	100	40	86-	all	EMP 1982								
Oct 14	13	01-16 Amilia	12.7	2.14	128970	9.0	G5 0 47.1	-3 18 3.7	11 23 22	Hawaii?s; eastern Indonesia	166	62	82-	e120°E	EMP 1982							
Oct 14	20	26-35 Winchester	11.1	1.57	11.2	6 35.7	2 32 0.7	17 20 11	Mauritius?n; southwest Australia	101	33	79-	all	EMP 1982								
Oct 15	15	24-29 Winchester	11.1	1.56	9.5	88	6 36.5	2 29 1.8	17 20 11	northern Australia; New Zealand?s	101	27	73-	all	EMP 1982							
Oct 16	0	29-51 Marghanna	11.7	1.04	11.4	3 07.0	13 03 0.9	8 23 20	M'ritius,Africa,Canaries;B'muda?n	154	42	69-	e 50°W	EMP 1983								
Oct 17	14	47-50 Penelope	12.2	1.95	163019	9.2	K2 19 46.9	-19 29 3.0	5 14 31	south central U. S. S. R.	91	173	53-	none	EMP 1981							
Oct 17	16	17-23 Winchester	11.1	1.54	11.7	6 38.5	2 21 0.5	18 21 11	eastern Siberia, Alaska	103	27	52-	all	EMP 1982								
Oct 19	12	00-10 Winchester	11.0	1.53	12.6	6 40.2	2 14 0.2	19 23 11	Hawaii?n; western Mexico?n	104	44	32-	all	EMP 1982								
Oct 20	12	24-29 Winchester	11.0	1.52	10.5	6 41.1	2 10 1.1	20 23 11	Alaska, western Canada	105	56	22-	all	EMP 1982								
Oct 21	4	02-26 Messalina	14.4	2.84	57899	8.5	K2 5 16.3	36 53 5.9	17 53 39	Libya,Sicily,France,J.K.,n.Canada	125	79	16-	e 55°W	EMP 1982							
Oct 21	9	35-60 Iris	8.4	1.14	12.0	5 47.3	26 28 0.0	42 39 7	Hawaii, Mexico, Jamaica	121	77	14-	e105°W	Branham								
Oct 21	14	53-57 Pretoria	12.9	2.34	73865	9.5	A5 0 14.2	27 16 3.5	13 22 19	western U. S. S. R.	153	141	12-	none	EMP 1981							
Oct 23	2	15-21 Winchester	11.0	1.50	13.1	6 43.3	2 01 0.1	22 26 10	United Kingdom, Europe	107	89	3-	e 35°E	EMP 1982								
Oct 24	21	05-23 Boliviana	11.3	1.30	108591	8.8	M0 23 21.5	10 53 2.6	17 30 15	Newfoundland?w;Azores?e.e. Brazil	142	137	0+	none	EMP 1982							
Oct 25	23	52-58 Hebe	9.3	1.55	11.4	6 45.2	4 47 0.1	26 34 12	South Africa	109	129	3+	none	Branham								
Oct 31	6	58-62 Eugenia	11.5	1.95	12.0	1 58.0	2 32 0.5	19 21 11	northern Canada, Alaska	167	83	48+	w125°W	Hergert77								
Nov 2	7	26-50 Winchester	10.9	1.42	9.8	6 49.9	1 30 1.4	38 43 10	Mexico?n; northern South America	114	127	67+	w 90°W	EMP 1982								
Nov 2	20	31-44 Doris	11.1	1.98	11.5	2 21.6	8 18 0.6	16 22 14	cen. & sw USSR, Turkey, nw Africa	173	60	72+	w 60°E	Hergert77								
Nov 4	0	04-27 Winchester	10.8	1.41	10.7	6 50.6	1 26 0.8	43 49 10	U. K., s. Europe, Mideast, India	116	110	81+	w 30°E	EMP 1982								
Nov 4	12	46-57 Hermione	13.2	3.19	10.6	8 19.9	24 01 2.7	22 38 23	Hawaii?s; northwest U. S. A.	100	126	85+	w150°W	EMP 1983								
Nov 6	7	05-18 Eugenia	11.6	1.97	12.4	1 53.2	2 06 0.4	21 23 11	northwest South America; Tahiti?s	161	12	95+	all	Hergert77								
Nov 6	12	01-27 Winchester	10.8	1.39	11.3	6 51.6	1 21 0.5	54 61 10	Hawaii; western Mexico?n	118	84	95+	w115°W	EMP 1982								
Nov 8	5	22-30 Pretoria	13.1	2.46	10.8	F5 0 07.3	24 26 2.4	14 25 20	e.U.S.A.?n; Hispaniola, Panama	141	36	100+	all	EMP 1981								
Nov 12	0	04-25 Hypatia	11.9	1.79	112399	7.6	G0 5 02.4	6 04 4.3	16 26 17	Mideast,central Africa, s. Brazil	150	22	90-	e 45°W	EMP 1983							
Nov 12	12	58-71 Themis	11.6	2.19	13.2	2 31.1	15 02 0.2	17 22 14	Hawaii?n; Indonesia	170	53	86-	e120°E	Hergert78								
Nov 13	4	35-51 Ceres	7.3	1.83	10.3	3 08.7	8 39 0.1	79 21 3	nwAfrica, Caribbean,Mexico,Hawaii	171	56	81-	e120°W	APENAXX								
Nov 15	11	46-48 Euphrosyne	11.7	2.64	10.1	11 31.1	29 54 1.8	10 12 14	Mexico?n	74	33	59-	all	Hergert78								
Nov 19	1	34-36 Venus	-4.6	1.14	186841	2.8	K0 18 24.9	-25 27 0.0	296 5 1	southern Alaska, British Columbia	39	92	20-	none	NA0001							
Nov 22	9	34-36 Euphrosyne	11.7	2.58	11.9	11 41.2	29 27 0.6	11 13 14	northern Canada, Greenland	78	71	0-	none	Hergert78								
Nov 23	7	00-05 P/Halley	21.3	4.81	12.9	6 24.9	11 59 8.5	3 15 70	Canary Islands, Florida, Mexico	143	147	0+	none	Yeomans7								
Nov 24	9	23-40 Faina	11.4	1.28	11.1	G5 4 48.8	17 22 0.9	10 20 16	Venezuela, Mexico, Japan	168	169	3+	none	EMP 1982								

Table 1, Part E

1984 DATE	UNIVERSAL TIME	P NAME	L NAME	A NAME	N NAME	E NAME	T NAME	S NAME	SAO NAME	A NAME	R NAME	O NAME	C NAME	U NAME	L NAME	T NAME	A NAME	T NAME	I NAME	O NAME	N NAME	Ephem. Source		
					<u>m_v</u>	<u>Δ_{AU}</u>	<u>Δ_{AU}</u>	<u>SAO No</u>	<u>m_v</u>	<u>Sp</u>	<u>R.A. (1950)</u>	<u>Dec.</u>	<u>Δm</u>	<u>Dur</u>	<u>df</u>	<u>P</u>	<u>Possible Area</u>	<u>SUN</u>	<u>El</u>	<u>M</u>	<u>0</u>	<u>0</u>	<u>N</u>	<u>Up</u>
Nov 24	12 45-47	P/Halley	21.3	4.78	13.0	6 ^h 23 ^m	11°58'8.3"	3°15'69"	New Zealand?s; Papua	144°161°	3+	none	Yeomans7											
Nov 24	23 57-62	P/Halley	21.3	4.77	12.9	6 23.5	11 58 8.4	3 15 69	India?n; central Africa	145 164	5+	none	Yeomans7											
Nov 28	5 52-87	Winchester	10.5	1.26	114569.0	A3	6 50.9	1 27 1.7	49 52	9 Brazil, Ecuador, see note	137 141	30+	none	EMP 1982										
Dec 1	15 24-50	Iris	7.3	0.93	11.5	5 39.1	23 50 0.0	34 29	6 Alaska?n; e.Siberia, China, India	164 92	62+	w135°E	Branham											
Dec 3	22 26-39	Harmonia	10.0	1.36	78926	9.2	6 55.5	22 19 1.3	17 33	17 central Africa	149 82	82+	all	Herget78										
Dec 7	13 48-75	Iris	7.2	0.93	11.1	5 33.2	23 18 0.0	31 27	6 w. U.S.A., Japan, China, e. India	171 19	99+	all	Branham											
Dec 9	4 40-58	Winchester	10.3	1.22	12.3	6 44.8	2 16 0.2	28 29	9 northern South America; Mexico?n	148 28	99-	all	Herget78											
Dec 10	23 15-18	Berbericia	13.5	3.40	100286.3	F8	12 46.8	14 03 4.3	8 15 27	Arabia, India, south central China	74 75	93-	all	Herget78										
Dec 11	23 16-35	Winchester	10.3	1.21	12.3	6 42.8	2 34 0.2	25 27	8 South Africa, eastern Brazil	150 33	87-	e 45°W	Herget78											
Dec 12	8 46-52	Winchester	10.3	1.21	12.6	6 42.5	2 37 0.1	25 26	8 eastern Canada?s	151 36	83-	all	EMP 1982											
Dec 12	9 53-69	Hermione	12.8	2.75	12.9	8 21.0	25 32 0.7	23 37	20 Colombia, Mexico; Hawaii?s	138 7	83-	all	EMP 1983											
Dec 13	7 51-70	Iris	7.0	0.93	11.4	5 27.1	22 46 0.0	30 26	6 northern U. S. A.; Hawaii?s	179 59	75-	e155°W	Branham											
Dec 14	11 18-45	Iris	7.0	0.93	9.9	5 25.9	22 40 0.1	30 26	6 Mexico,Hawaii?n;Papua,w.Australia	179 75	63-	e175°E	Branham											
Dec 15	16 05-18	Tercidina	12.5	1.72	109396	8.9	F8	0 40.1	3 31 3.6	11 27 23	Kenya; Sri Lanka?n; Malaysia	107 163	50-	e115°E	EMP 1981									
Dec 16	4 12-29	Hebe	8.5	1.26	11.7	6 29.9	4 02 0.1	19 22	10 nw Africa, ne U. S. A., sw Canada	156 85	44-	e 60°W	Branham											
Dec 16	22 29-44	Dejopeja	13.1	2.22	10.0	C44	5 31.4	24 49 3.1	10 21 24	S'mtra,India,sArabia,nAfr,Am'znR	178 108	35-	e 40°E	EMP 1983										
Dec 17	0 48-60	Athor	12.5	1.65	59154	8.9	G0	6 27.0	37 05 3.6	8 20 24	Sinkiang, Kazak SSR, Urals, northern and western Canada, northwest USA	163 97	34-	e 15°E	EMP 1981									
Dec 20	4 50-74	Iris	7.2	0.95	11.1	5 20.0	22 08 0.0	31 27	6 northwest Africa, Florida, Mexico	173 158	7-	e 2°W	Branham											
Dec 22	3 41-59	Winchester	10.2	1.21	11.5	6 34.1	3 59 0.3	20 21	8 w. cen. Africa, ne USA, sw Canada	159 155	0-	none	EMP 1982											
Dec 23	2 21-28	Winchester	10.2	1.21	12.9	6 33.2	4 09 0.1	20 21	8 northern South America; Mexico?n	159 157	1+	none	EMP 1982											
Dec 23	4 12-26	Winchester	10.2	1.21	11.3	6 33.2	4 10 0.3	20 21	8 northern South America; Mexico?n	160 157	1+	none	Herget78											
Dec 25	23 51-68	Winchester	10.2	1.21	9.3	A0	6 30.5	4 39 1.3	20 20	8 s and w Africa, Georgia (U.S.A.)	161 135	14+	w 55°W	EMP 1982										
Dec 27	1 10-18	Winchester	10.2	1.21	11.3	6 29.6	4 50 0.3	19 20	8 Iran, western U. S. S. R.	161 123	22+	none	EMP 1982											
Dec 27	14 30-35	P/Halley	20.9	4.36	12.4	5 50.4	12 00 8.5	3 11 63	Hawaii, China, India	166 109	26+	w110°E	Yeomans7											
Dec 29	4 56-59	Eugenia	12.4	2.51	10.3	1 39.1	2 16 2.2	33 41	15 northwest South America	106 28	40+	w 83°W	Herget77											
Dec 30	19 00-19	Ate	10.8	1.36	78743	9.5	F8	6 44.3	26 48 1.5	16 23 13	Indonesia, Sri Lanka, cen. Africa	176 85	55+	w 80°E	EMP 1979									
Dec 30	20 08-13	Winchester	10.2	1.22	10.6	6 26.1	5 33 0.6	19 20	9 South Africa?n	162 81	56+	all	Herget78											
Dec 31	5 00-01	Nuwa	12.5	2.04	96300	9.3	A5	6 54.0	19 43 3.2	10 21 22	Patagonia	175 83	59+	w 69°W	EMP 1980									

Neptune is from "Predicted Occultations by Neptune, 1981-1984" by D. Mink, A. Klemola, and J. Elliot, *Astronomical Journal* 86 (12), 135. That article and another one by the same authors in the same issue of *Astron. J.* list some occultations of fainter stars by Uranus and by Neptune, which probably will be detectable only with photoelectric equipment used with large telescopes at (Text continues on next page)

Table 3.

Ephemeris Differences for 1984					Ephemeris Source
Date	MP#	Shift	At		
Jan 12	154	1°24S	10 ^m .2		Herget 78
Jan 14	49	0.30S	1.1		EMP 1983
Jan 16	154	1.38S	7.5		Herget 78
Jan 20	165 (both)	0.06S	0.2		EMP 1980
Jan 26	41	0.45N	3.5		EMP 1976
Jan 29	46	0.02S	-0.7		EMP 1980
Jan 31	154	1.51S	3.3		Herget 78
Feb 10	154	1.49S	2.5		Herget 78
Feb 22	268	42.75S	-63.0		Herget 78
Feb 23	386	0.93S	-1.7		Herget 77
Feb 25	31	2.87S	8.7		Herget 78
Mar 4	241	0.59N	-11.2		Bardwell 67
Mar 23	114	0.24N	1.7		EMP 1982
Mar 26	10	0.09S	-4.8		Klepczynski
Mar 31	11	1.23S	5.0		EMP 1980
May 2	128	0.44S	3.2		EMP 1977
May 11	230	1.87N	5.1		Herget 78
May 19	230	2.03N	5.7		Herget 78
May 30	104	0.02N	68.8		EMP 1980
Jun 1	85	0.06S	-1.4		ITA 1977
Jun 2	47	0.06N	1.0		EMP 1983
Jun 2	602	1.25S	-21.0		EMP 1981
Jun 2	624	0.68N	-56.4		Herget 78
Jun 4	70	0.60E	0.5		EMP 1980
Jun 8	13	0.07N	-0.9		EMP 1980
Jun 24	145	0.61N	1.3		Herget 78
Jul 14	139	0.36S	6.9		ITA 1974
Jul 16	386	1.78S	-0.1		Herget 77
Jul 16	211	0.25S	-0.1		EMP 1983
Aug 8	87	0.12N	-1.8		EMP 1983
Aug 19	747	1.70N	-3.3		Herget 78
Aug 21	747	1.74N	-3.4		Herget 78
Aug 25	104	1.62S	-167.3		EMP 1980
Aug 28	747	1.89N	-3.5		Herget 78
Sep 2	747	2.00N	-3.6		Herget 78
Sep 16	47	0.06S	-2.3		EMP 1983
Sep 21	747	2.46N	-4.4		Herget 78
Sep 23	747	2.53N	-4.5		Herget 78
Sep 23	747	2.54N	-4.5		Herget 78
Oct 2	334	0.94S	39.2		Herget 78
Oct 12	804	0.56S	-13.2		EMP 1982
Oct 14	747	3.17N	-6.7		Herget 78
Oct 15	747	3.20N	-6.9		Herget 78
Oct 21	790	0.55N	-3.0		Herget 78
Oct 24	712	0.16W	-0.7		Herget 78
Nov 2	747	3.84N	-14.7		Herget 78
Nov 6	747	4.00N	-20.8		Herget 78
Nov 8	790	1.01N	-2.6		Herget 78
Nov 28	747	4.85N	18.3		Herget 78
Dec 3	40	0.91N	6.5		EMP 1975
Dec 10	776	4.13S	-17.5		EMP 1980
Dec 11	747	5.43S	-9.0		EMP 1982
Dec 16	184	0.50N	2.6		Herget 78
Dec 23	747	5.83N	6.8		Herget 78
Dec 25	747	5.90N	6.3		Herget 78

which will occur before the next issue of *O.N.* is distributed, I will try to distribute advance predictions to regional coordinators.

History of Astrographic Catalog Searches. IOTA obtained magnetic tape copies of several of the northern Astrographic Catalog zones from the Stellar Data Center in Strasbourg, France, in 1977, as described in *O.N.* 1 (9), 97. They were obtained to include fainter stars in predictions for both lunar and asteroidal occultations. The data were a little difficult to use, being ordered by the astrographic plates, rather than in right ascension order. I wrote a computer program to extract data for selected small regions and sort the stars within the region in order of increasing right ascension. This was used mainly for star clusters to form the main basis for the USNO "J" and later "C" catalogs for extended lunar occultation predictions, first distributed in late 1977, as described in *O.N.* 1 (12), 123. Occultations of A.C. stars by asteroids were first computed at the same time and published in *O.N.* 1 (12), 127. But the latter work was haphazard, depending on manual searches by D. Wallentinsen and others to identify when an asteroid would cross an open cluster, so relatively few events were published. The relatively high frequency of A.C. events for a given asteroid was demonstrated with my articles, "Occultations of A.C. Stars by (24) Themis," in *O.N.* 2 (5), 43 and "Occultations of A.C. Stars by (27) Euterpe" in *O.N.* 2 (7), 73. In spite of the early 1900's epoch of most of the A.C. plates, modern astrometry by Klemola and Penhallow has demonstrated that the A.C. positions are within 1" of the current positions for the large majority of faint A.C. stars, an accuracy which rivals that of most asteroid ephemerides. Most of these faint stars are very distant and have negligible proper motions. However, when possible, new astrometric observations are recommended to improve the A.C. position and identify stars whose positions now differ from the predicted positions due to significant proper motion.

For a long time, I had planned to compare asteroid ephemerides with A.C. data automatically to find occultations of many stars not in the AGK3 and SAO catalogs. But other commitments prevented my making the important first step of sorting all of the A.C. data from France into right ascension order and eliminating duplicates (most A.C. stars appear on at least two plates). Alain Fresneau finally accomplished this job at the Space Telescope Science Institute in Baltimore last year, to help set up the Guide Star Selection System for the Space Telescope. He published his procedures in an article, "Survey of the Astrographic Catalog from 1 to 31 Degrees of Northern Declination," in *Astron. J.* 88 (9), 1378 and gave a formatted magnetic tape with 1,025,208 A.C. stars to Wayne Warren at the Astronomical Data Center at Goddard Space Flight Center for distribution to the astronomical community (A few of the stars on Fresneau's tape were out of order; Warren corrected this). I obtained a copy and produced a binary version at USNO for more efficient use with the IBM 4341 computer there. During the course of this work, I found 27,897 duplicate entries, less than 30 of which I felt were possible true double stars. I deleted all of the duplicates from my binary copy, but sent a tape containing only the duplicates to Fresneau, so that he could trace their origin. I modified my occultation search computer

program to read the binary tape for comparison with both retrograde and prograde parts of an ephemeris. The program would not find occultations which occur within a couple of days of a stationary point or near the 24h - 0h discontinuity in right ascension, but stars near these troublesome points were printed so that any events missed in the automatic part of the search could be found manually.

Occultation of Astrographic Catalog Stars during 1984. Since some manual work still was involved and time was short for including the A.C. events with the others for 1984 in this issue of *O.N.* (the work was begun in mid-November, 1983, it was not practical to consider the hundreds of asteroids involved in the searches performed at the Royal Greenwich and Lowell Observatories. I limited my search to those asteroids with diameters of at least 200 km (according to Tedesco, see above) which have oppositions between declinations of -2° and $+33^\circ$ during 1984 or late 1983, as well as a few smaller asteroids of special interest. Unfortunately, (532) Herculina, (171) Ophelia, and several other interesting asteroids are in the Southern Hemisphere, entirely south of the area covered by the French A.C. The asteroids which I included in my searches for 1984 have the following numbers: 1-4, 6, 7, 9 (occultation binary suspect), 12 (speckle binary suspect), 19, 24, 31, 41, 45, 48, 49, 65, 88, 121, 130, 154, 165, 194, 324, 375, 386, 423, 624, 747, and 2060. Halley's Comet (P/Halley) was also included; its ephemeris was generated from the latest orbital elements supplied by Don Yeomans at J.P.L. and used a numerical integration program which includes non-gravitational forces. The ephemerides which I generated for the searches were restricted to the declinations noted above and to elongations greater than 45° from the sun. The faint A.C. stars would be difficult to observe closer to the sun (favorable closer-elongation events of brighter SAO and AGK3 stars presumably would have been found by Taylor). Occultations also were rejected if the asteroid angular diameter was less than $0''.08$, when even last-minute astrometry likely would give a poor prediction of the path relative to its width. This restriction was waived in the case of asteroids of special interest, such as (9) Metis, (12) Victoria, (624) Hektor, and (2060) Chiron. No Chiron events were found. Both Cincinnati (Herget) and Leningrad ephemerides were used for the searches for (154) Bertha and (747) Winchester, since these differed by two or more arc seconds and recent observations could not distinguish which is the better ephemeris (in each case, the Leningrad orbit seems to be favored slightly). For the others, I have used what I consider to be the best orbit, the source being given in Table 1 for individual events.

Observational Considerations for the Astrographic Catalog Occultations. Many of the asteroids considered in my A.C. searches are considerably brighter than the faint A.C. stars, so that their occultations can be recorded only photoelectrically. This is especially the case for (1) Ceres and (4) Vesta; no prediction is given if the blue-magnitude Δm is less than 0.05. However, the Δm 's in the table are computed from asteroid V magnitudes, while only photographic (blue) magnitudes are available for the A.C. stars. This is for conformance with other catalogs, where V or photovisual magnitudes are available, and a visual Δm is desired. But most faint A.C. stars are brighter visually (because most have

later spectral types than A0) than their A.C. magnitude, sometimes by more than a magnitude. Hence, the actual Δm 's often will be a few to several tenths larger than given in Table 1 for the A.C. stars. So visual observers should check the relative magnitudes of the star and asteroid if the predicted Δm is about 0.4 or larger.

Finder charts have been prepared with an expanded scale from the A.C. data for a square 1 degree on a side centered near the target star. This square is shown in the AGK3-based plot drawn at the same scale as finder charts appearing in previous issues of *O.N.*, which, however, are now only 3° on a side to conserve space. In some cases, small-scale Atlas-Coeli-based charts are not included due to lack of time. A.C.-based charts also are included for AGK3 stars fainter than about mag. 8.5, since the fainter stars help to identify the proper target star. Since Ceres and Vesta are brighter than 8th magnitude, they can be located reliably with the AGK3-based charts. Therefore, A.C.-based charts usually are not produced for them. In general, only the A.C.-based charts in *O.N.* have been checked against the True Visual Star Atlas for Northern Hemisphere stars. I do not do this manual checking for unpublished finder charts which I distribute directly to regional coordinators in other countries.

When Fresneau produced his tape, none of the A.C. designations were saved. Hence, no numbers are given for these stars; they can be identified only by their coordinates and magnitudes. Fresneau may save the original designations on a future version of his tape. In the meantime, we will use the current tape as well as we can. When Lowell and the Royal Greenwich observatories obtain copies of Fresneau's tape, they probably will use their more efficient (at least, more automatic) computer-search programs to extend the A.C. searches to smaller asteroids. My programs also might be used on other computers by one or two other IOTA members to extend the searches, or I will do it if I can find time to automate some of the manual procedures currently used. I also would like to add more catalog data, especially Yale Catalog stars which are in neither the AGK3, SAO, or Perth 70 catalogs, and IOTA's Southern Astrophotographic Catalog Project stars. Extension to all asteroids larger than 0.08 will result in many hundreds, even thousands, of events per year, and current publication methods will not be practical. An automatic method of geographically sorting the predictions will be needed, so that observers would be sent computer-produced data only for events in their regions, as we do now for lunar occultations. Then, only the very best events would be included in *O.N.*, mainly for the benefit of those who might travel to other regions for outstanding occultations. I have plans for how this might be accomplished, but it will take time to implement them.

Ephemerides and Stellar Data. General guidelines for the ephemerides used for my predictions were given in my article about 1983 events referenced above. Table 3, listing ephemeris differences for several 1984 events, is in the same format as last year's table. My comparisons with recent astrometric observations, aided by Brian Marsden and Conrad Bardwell at the Minor Planet Center in Cambridge, MA, showed that the Leningrad (Ephemerides of Minor Planets, or EMP) orbits were slightly better than Herget's orbits for (31) Euphrosyne, (241) Germania,

(624) Hektor, and (747) Winchester. However, the differences were not very significant except for (624) Hektor. (194) Prokne's orbit was rated poor by Millis *et al.*, so they excluded it from their searches (Herget did not compute an orbit for Prokne). However, I found that two of the four pairs of observations made since 1976 agreed within about 1" with the positions computed from the EMP 1983 orbit. On closer examination, it was found that the other two pairs would also be in good agreement if the observers had made a 1-hour error in the time of their exposures. An observation in November by Klemola confirms the good accuracy of the Leningrad orbital elements. On the other hand, the orbits for (268) Adorea disagree with recent data by tens of arc seconds for Leningrad and by several arc seconds for Herget. Herget's orbit, apparently the better one, predicts a wide miss, so the February 22nd occultation probably will not happen. The orbit for (365) Corduba is also poor, so the occultation predicted by it on October 6 probably will not occur.

I have used AGK3 positions and proper motions for many of my nominal predictions, rather than SAO data. This agrees with Taylor's and Lowell Observatory's practice, but it is the opposite of my practice in previous years. Good-quality astrometric observations usually favor the AGK3 data, but not always. Wayne Warren supplied me with Yale Catalog data for several stars, as well as some useful double star data. The individual components of four close double stars are listed separately in Tables 1 and 2.

Prediction Updates. The predictions for asteroidal occultations can be improved by astrometric observations to update the ephemeris and the star's position a few months in advance. The improvement can be quite good if the asteroid happens to pass near the star, perhaps during its retrograde loop, so that both objects can be photographed on the same plate. Sometimes these preliminary prediction improvements are published in *O.N.* or in *Sky and Telescope*, or distributed by mail by Gordon Taylor, by Lowell Observatory astronomers, or by me. But a very accurate prediction generally can not be made until the objects are close enough together to photograph on the same astrometric plate during their final approach, only a few days before the event. In these cases, those of us involved with computing the final predictions must concentrate on notifying those near the path; this is facilitated if those we contact telephone other observers in their areas. You can find out the latest predicted shift and time correction for upcoming asteroidal occultations by telephoning Astro-Alert in Chicago, IL, area 312, 259-2376, or Gordon Taylor at the Royal Greenwich Observatory, England, city code 0323, 833171, ext. 3252. Information also usually can be obtained from Paul Maley in Houston, TX, 713,488-6871 and, at least for events in the western U.S.A., from Lowell Observatory, Flagstaff, AZ, 602,774-3358.

Notes about Individual Events. There will be considerable moonlight interference for many of the 1984 events, especially for some of the faint A.C. stars, so pay attention to the lunar elongation and percent sunlit given in Table 1. For these cases, it would be useful to try to locate the star about a month before the occultation, when moonlight conditions might approximate those for the actual event. Any prior experience in locating the star, even in a

dark sky, will help on the night of the occultation. Notes about some of the events are given below:

Jan. 7, Halley's Comet: The nucleus is less than 100 km in diameter, but there may be some material in orbit 50 km or more from the nucleus.

Jan. 12, Halley's Comet: Klemola has measured a recent Lick Observatory plate to improve the star's position. His position is south of the A.C. position, which means that the occultation path is expected to shift north, off the earth's surface. But the ephemeris could be in error by at least 1", larger than the shift.

Jan. 20, Prokne: In early November, Prokne passed close to the star. Klemola obtained a plate, measurements of which indicate a 0:26 south shift and a correction to the time of 2.4 minutes early. The shift puts the path across Oregon, Louisiana, and southern Florida. Although the path is very likely to cross the U.S.A., the seeing was not good when the Lick plate was taken. This, combined with ephemeris drift, could cause the path to shift 0:5 from the above value.

Jan. 26: The star is Aitken Double Star (ADS) 3707. The 11.8-mag. B-component is 1:3 away, in position angle 344°. Good seeing will be needed in order to resolve the star and detect its occultation.




Diagram by Gordon Taylor
OCCULTATION OF SAO 158913 BY SATURN ON 1984 MARCH 25

Jan. 27: The note for Jan. 12 is also true for this event.

Feb. 25: The star is ADS 1654. The 8.6-mag. primary is 2:0 away in p.a. 344°; its occultation path misses the earth above Antarctica.

Mar. 4: The star is α Scorpii (Al Niyat or Zodiaca Catalog No. 2349 or ADS 10009), the brightest known to be occulted by an asteroid this year. A star brighter than Sigma is occulted by an asteroid larger than Germania about once every 35 years, on the average. The 3.0-mag. primary is probably a spectroscopic binary (hence, the double star code U) which might be resolved during this spectacular occultation. Lunar occultation observations show a 5.2-mag. component (component "2") which is probably now about 0:7 away, according to an analysis by Ed Nather and P. Wild given in *Publications of the Astronomical Society of the Pacific*, Vol. 86, p. 116. Hence, the Δm that actually will be seen when the primary is occulted will be 2.2, not 10.0, since the 5.2-mag. star will remain visible. It will be occulted along a separate path two minutes earlier and probably north of the primary, but the Δm will be only 0.1, detectible only photoelectrically, since the primary will remain visible. The 8.3-mag. visual B component, 20" away in p.a. 273°, will not be occulted from the earth, missing it by 2". I have made tentative plans to observe the occultation of the primary star. NOAA satellite data for early March during the last 3 years indicate relatively clear skies over southern China, Indochina, and the northern Philippines, but poor weather prospects for northern China and Japan.

Mar. 25: This occultation will be difficult to see due to the faintness of the star relative to Saturn, and even photoelectric attempts may fail. The star will be visible for several minutes after it reappears from behind the ball of the planet and before it merges into the inner edge of Ring C.

Apr. 10: The 10.3-mag. B component is 0:48 away in p.a. 63°.

Apr. 19, Tamara: SAO 207516 is the secondary of a double star. The 8.6-mag. primary, SAO 207517, is 4:9 away in p.a. 118°, according to the Lick Index Catalog of Double Stars. But the difference in the current SAO positions is about 8" and implies that the occultation path for the primary misses the earth by 2" above South America. The SAO used relatively poor G.C. data.

Apr. 24: The star is in M35. Fred Schaaf, Millville, NJ, informed me that this M35 passage had been predicted by Steve Albers.

May 2: The star is ADS 8938. Two 12th-magnitude companions 16" and 140" away will not be occulted.

May 12, Mars: The star is Z.C. 2173. This event will be very difficult, since Mars will be at opposition, with essentially no defect of illumination (dark crescent). Taylor predicts immersion at 15:23.0 at Tokyo, with emersion 17.0 minutes later in p.a. 78°. At Naini Tal, India, the corresponding times are 15:36.2 and 15:54.4 in p.a. 103°.


June 24: The star is 83 Leonis (Z.C. 1660 or ADS 8162). The 7.6-mag. B component (SAO 118865) is 28"

away in p.a. 149", while a 10th-mag. companion is 90" away. Neither will be occulted by Fortuna.


July 22: This occultation might be seen by using high power to spread out Neptune's image, so that a higher effective Δm can be achieved. The occultation may last less than 10 minutes for most Northern Hemisphere observers. The nominal northern limit passes just above the earth's north polar regions. The position I used combined SAO proper motion with Klemola's position. If Klemola's position alone were used, the northern limit of the occultation would cross North America.

Nov. 13: A possible north shift could bring more observatories in the southern U.S.A. into the path of this important event. A.C. data predict a path across the northern U.S.A., but the Klemola position for the star, used for the nominal prediction, is certainly better.


Nov. 19: The star is λ Sagittarii (Kaus Borealis or Z.C. 2672). Lunar occultations show that the star is single, with a diameter of 0".005. Venus will be 75% sunlit with a 3% defect of illumination. The location of the southern limit could be in error by 100 km or more. Central graze will be only 2° from the south cusp on the dark side as seen from the southern limit.




Use a tracing of this cosine scale to estimate star altitude from the Soma world maps; place the 90° mark at the center of the circle, and read the star altitude at the observing site.






EMERIS SOURCE - EMP 1983






SAO 118733 by Hestia 1984 Jan 29




Anonymous by Bertha 1984 Jan 29




Anonymous by Prokne 1984 Jan 30


1984 JAN 23 (121) HERMIONE
DIAMETER 201 KM - 0.001



1984 JAN 23 (121) HERMIONE
DIAMETER 201 KM - 0.001




EPHEMERIS SOURCE = EPH 1983




This plot showing the region of the head of Hydra is based on ATLAS ECLIPTICALIS. The arrow, in the approximate direction of (194) Prokne's motion, shows SAO 117122, to be occulted on 1984 Jan. 20. Four stars indicated by dots are shown on the TRUE VISUAL MAGNITUDE ATLAS but not on ATLAS ECLIPTICALIS. They are each almost as bright as SAO 117122, except for the star labelled A, which is 0.5 to 1.0 magnitude brighter than SAO 117122.


1984 JAN 23 (121) HERMIONE
DIAMETER 201 KM - 0.001




EPHEMERIS SOURCE = HERMIONE




Anonymous by Bertha 1984 Jan 31




Anonymous by Bamberg 1984 Feb 3




+15°2320 by Dembowska 1984 Feb 6




Anonymous by Bertha 1984 Feb 10




Anonymous by Prokne 1984 Feb 13




Anonymous by Bamberga 1984 Feb 14




Anonymous by Victoria 1984 Feb 15




SAO 186422 by Interamnia 84 Feb 17




Anonymous by Victoria 1984 Feb 17




SAO 119464 by Metis 1984 Feb 19




Anonymous by Prokne 1983 Feb 19




Anonymous by Prokne 1984 Feb 20






Anonymous by Victoria 1984 Feb 21




+16°1989 by Adorea 1984 Feb 22








Anonymous by Vesta 1984 Feb 23




SAO 141366 By Siegena 1984 Feb 23




Anonymous by Vesta 1984 Feb 24



-24° 30'28 by Euphrosyne 184 Feb 25




Anonymous by Prokne 1984 Feb 25





Anonymous by Metis 1984 Feb 25

1984 FEB 14 324 BAMBURGA


DIAMETER 205 KM - 0.003




EPYMERIS SOURCE - 1984 FEB 25





SAO 80634 by Andromache '84 Feb 26




Anonymous by Metis 1984 Feb 28




Anonymous by Prokne 1984 Mar 1



Anonymous by Vesta 1984 Mar 2



-25° 11485U by Germania 1984 Mar 4



Anonymous by Elektra 1984 Mar 5

1984 MAR 5 10:00 AM
1984 MAR 5 10:00 AM




1984 MAR 5 10:00 AM
1984 MAR 5 10:00 AM


1984 MAR 5 10:00 AM
1984 MAR 5 10:00 AM




1984 MAR 5 10:00 AM
1984 MAR 5 10:00 AM




Anonymous by Barberga 1984 Mar 5




Anonymous by Euphrosyne 1984 Mar 5




Anonymous by Thisbe 1984 Mar 10




Anonymous by Victoria 1984 Mar 10




SAO 118750 by Euterpe 1984 Mar 13




Anonymous by P/Halley 1984 Mar 14




Anonymous by Metis 1984 Mar 15



Anonymous by June 1984 Mar 16




Anonymous by Prokne 1984 Mar 16




SAO 159989 by Kassandra '84 Mar 23

1984 - 4 (241) ALMANAC SAO 104336
 DIAMETER 187 KM 0.038



EPHEMERIS SOURCE - EMP 1981

1984 - 4 (241) ALMANAC SAO 110150
 DIAMETER 110 KM 0.013



EPHEMERIS SOURCE - 186717

FIGURE 1. GALACTIC COORDINATES
TANGENT GALACTIC PLANE



FIGURE 2. GALACTIC COORDINATES
TANGENT GALACTIC PLANE










FIGURE 3. GALACTIC COORDINATES
TANGENT GALACTIC PLANE

EMERIS SOURCE = EMP 1983

FIGURE 4. GALACTIC COORDINATES
TANGENT GALACTIC PLANE


1984 3 24 (114) KASSANDRA SAO 159389
DIAMETER 143 KM - 5.03

1984 3 24 (114) KASSANDRA SAO 159389
DIAMETER 143 KM - 5.03






1984 3 24 (114) KASSANDRA
DIAMETER 143 KM - 5.03

EPHEMERIS SOURCE - HERGET78



EPHEMERIS SOURCE - HERGET78


EPHEMERIS SOURCE - HERGET78




SAO 158913 by Saturn 1984 Mar 25

Anonymous by Juno 1984 Mar 26


SAO 163443 by Hygiea 1984 Mar 26




SAO 162170 by Penelope 1984 Mar 27



16" 2571 by Parthenope 1984 Mar 31




Anonymous by Metis 1984 Apr 2




Anonymous by Bamberg 1984 Apr 3

1984 4 9 (12) VICTORIA
DIAMETER 133 KM = 0.007




EPHEMERIS SOURCE = HERGET77




1984 APR 9 (12) VICTORIA


1984 APR 9 (12) VICTORIA




1984 MAY 4 (12) VICTORIA




1984 MAY 3 (12) VICTORIA




1984 MAY 23 (12) VICTORIA



Anonymous by Victoria 1984 Apr 9



Anonymous by Ursula 1984 Apr 10



SAO 138680 by Meliboea 1984 Apr 10

CORRECTION CORRECTED

In order to hasten final assembly and mailing of this issue, the first twenty pages were taken to the printer before the figures for the remainder were available. In the interim, David Dunham phoned to say that the comment about the spelling of the name of asteroid (36) in the article "Planetary Occultations During 1984" is incorrect; the correct spelling is 'Atalante,' not Atlante sorry, the page number cannot be quoted, as the reference copy is still at the printer as this is being written.

We take this opportunity to mention that the tally coupon for 1983 total occultation timings is included in this issue, at least for those receiving the initial mailing.

The following article and notes are also last-minute additions to this issue.

THE INTERNAL REVENUE SERVICE MAKES IT OFFICIAL

On January 3rd, Paul Maley phoned to say that word at last has been received from the I.R.S. that IOTA's request for tax-exempt status has been granted. In case anyone needs to know, our tax number is 36-3254326. We understand, from Maley, that the tax-exempt status is retroactive, for some purposes, to August 18th, 1983. We hope that someone will provide us with a further report on the ramifications of this action, for our next issue.

Maley emphasized that participation in the IOTA expedition for the total solar eclipse of 1984 November 23 now may be considered to come under the category of charitable contributions, at least in part, and for qualified individuals. See *O.N.* 3 (5), 98 to 100.

SOME IMPORTANT NOTES ABOUT ASTEROIDAL OCCULTATIONS

David W. Dunham

With the larger number of events during 1984, no astrometric updates will be made for many of them, even many possible in North America. Observers should make a habit of monitoring all observable ap-pulses with miss distance less than 1"0 (typical error) at their location, or even within 2"0 (possible error for many events), when weather permits. Then, at least a few chords can be obtained and the asteroid's size determined if the occultation path does cross populous regions. No astrometry is likely for the occultation by (154) Bertha on Jan. 16 (a

conference on astrometry is being held at that time in Florida) and by (704) Interamnia on Feb. 17 (the star is relatively bright at mag. 8.0, but the altitude is too low to obtain astrometry in the Northern Hemisphere).

All of the finder charts in this issue have been checked with the True Visual Magnitude Atlas (TVMA). In general, the agreement with the A.C.-based charts is very good, even though blue-sensitive plates were used for the A.C. On the detailed chart for the occultation by (121) Hermione on January 23, two (probably red) stars (one, the star to be occulted, or "target star") are marked "FAINTER" because they appear much fainter on the TVMA, nearly as faint as the faintest A.C. stars shown. The A.C. plate magnitude limits are not uniform, and sometimes the target star does not appear in TVMA. These stars, indicated by notes on the charts, likely will require large telescopes for effective monitoring.

A good finder, such as an 8x50 with a field of view of at least 3°, is almost a necessity for locating objects in unusual parts of the sky, and is highly recommended. A very low-power, wide-field eyepiece (such as one with a 40-mm focal length) also is recommended with the main telescope for locating the target star. Once the star is identified positively, switch to a higher power if it is faint; this decreases the background brightness, which facilitates monitoring the star. For faint stars, use the highest power which still permits smooth tracking during the observing period.

Paul Maley's good article about last May's occultation of 1 Vulpeculae by (2) Pallas has been published in the current issue of *Astronomy* magazine.

NOTE: ADDITIONS TO THE MINUTES OF THE IOTA MEETING OF 1983 NOVEMBER 11

During the meeting, Paul Maley discussed plans for observing both solar eclipses in 1984; a graze of α Librae during the total lunar eclipse of May, 1985; and Halley's Comet and a lunar eclipse in early 1986.

Nine of the ballots designated D. W. Dunham as the proxy, and like Charles Herold, he cast all votes in favor of the motions.

The possible problems of the U. S. Naval Observatory are due to the departure of Dr. Van Flandern, rather than to budget cuts; see IOTA NEWS, p. 113.

D. W. D.